Plant Disease Reporter

Plant Science Research Division

Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 55 NUMBER 13

1971 Index
(Issued June 1972)

BIOLOGICAL SCIENCES
JUL 14 1972
LIBRARY

The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. Plant Science Research Division, in publishing this material, does not assume responsibility for the subject matter.

50:59
U5

PLANT DISEASE REPORTER INDEX

Issued by

PLANT SCIENCE RESEARCH DIVISION

Plant Industry Station
Volume 55

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABAWI, GEORGE S., 617, 1000
ABNEY, T. S., (776)
ADAIR, C. N., 864
ADAMS, P. B., 1077
AGRIOS, G. N., 869, 1049
AICHELE, MURIT D., (344)
ALCONERO, RODRIGO, 506, 902
ALEXANDER, ANN F., (929)
ALLEN, O. N., (145)
ALLEN, T. C., Jr., 546
ALLISON, ALLEN H., (419)
ANAGNOSTAKIS, SANDRA L., (199)
ANDERSON, GERALD W., 650
APPLEGATE, HART W., 625, 697
ARAGAKI, M., 392, (521), (743)
ARP, GREGORY, 577
ARTMAN, J. D., 834
AVERRE, C. W., (751)
AYERS, J. E., (401), (495)

BARGER, J. H., 271
BARKER, L. N., (820)
BARKSDALE, T. H., 253, 807
BARNES, GEORGE L., 711
BARNETT, O. W., 411
BARRITT, B. H., 135
BARTLETT, ALLAN B., (560)
BASSETT, M. J., (1061)
BEARD, J. B., (1118)
BECKETT, J. B., (401)
BEDI, P. S., (283)
BENOIT, A., (349)
BERGESON, G. B., 55
BERGQUIST, R. R., (392), 394, 941
BERRY, FREDERICK H., 1085
BETZ, JOE, (1136)
BHOWMIK, T. P., 87
BIEHN, W. L., 179, 968
BIRCHFIELD, W., 362, 1043
BITTENBENDER, H. C., (651)
BLACK, L. L., (599)
BLANCHARD, K. L., (726)
BLANPIED, G. D., (791)
BLODGETT, EARLE C., 344
BOCKHOLT, A. J., (202)
BOLLENBACHER, KATHARINA, 897
BOOTH, J. A., 569
BOPEARATCHY, R. N., (1021)
BORECKA, H., 828
BORECKI, Z., (828), 932
BORN, GERALD L., 996
BOUWKAMP, J. C., 1097
BOYD, R. J., 145
BRAVERMAN, SAMUEL W., 454, (639)
BRENDLER, R. A., (433)
BRISTER, L. R., (1043)
BROADBENT, P., 998
BROOKS, H. J., (379)
BROWDER, L. E., 319, (718)
BROWN, H. DANIEL, 317
BROWN, JOHN F., (885)
BRZESKI, M. W., 399
BUCHHOLTZ, W. F., (479)
BUFORD, W. R., (183)
BURKE, D. W., (1056)
BURTON, C. L., 1037
BUTCHER, CLYDE L., 54
BYSTYDZIENSKA, K., (932)
BYTHER, R. S., 7

CALDWELL, RODNEY W., (387)
CALLIHAN, R. H., (550)
CAMERON, H. R., 421
CAMPANA, R. J., (1085)
CAMPBELL, R. N., 322, (328)
CASE, F. W., Jr., (1108)
CHEE, K. H., 152
CHEW, C. F., (569)
CHIELDS, J. F. L., (241)
CHRISTIE, S. R., (1061)
CLARK, W. A., (871)
CLEMONS, G. P., 280
CLOSE, R., (45)
COAHRAN, D. R., (234)
COBB, F. W., Jr., (659)
COHEN, SHLOMO, (604)
COLE, H., Jr., (495), (684)
COLLINS, F. C,. (1053
CONSTANTINESCU, O., 733
CONVERSE, RICHARD H., (220), 560
COOK, A. A., (783), 785
COOK, J. A., (580)
COOPER, VALERIE C., (268)
CORDELL, C. E., 700
COVEY, RONALD P., Jr., 514
COX, E. A., (687)
COX, R. S., 350, 848
COYIER, DUANE L., 263
COYNE, D. P., 203, (505), (577)
CRAIG, JEWEUS, 672
CRAWFORD, JOHHNY L., 504
CRILL, PAT, 127
CROSIER, D. C., (483)
CSONKA, ELIZABETH, (77)

58549
U5

CZABATOR, F. J., 764

DAINES, ROBERT H., 746, 774
DAVE, G. S., (95)
DAVIS, J. R., 550
DAVIS, R. G., 1111
DAVISON, ARLEN D., 57
DEAN, LESLIE L., (54)
DeNEVE, R. T., (587), (851)
DICKENS, L. E., (609), 613
DIMOCK, A. W., (832)
DIMOND, A. E., (179)
DOOLEY, H. L., 1014
DRAKE, CHARLES R., 122
DRORI, TIRZAH, (431)
DuCHARME, E. P., (923)
DuRANT, J. A., 273
DURBIN, R. D., (361), (816)
DUTT, S., (283)
DYE, D. W., (517)

EBENEBE, Ch., 691
EDWARDS, D. I., 114, 415, 927, 974
EL-BEHADLI, A. H., (192)
ELGIN, J. H., 621
ELLETT, C. WAYNE, (382)
EL-NIL, M. M. ABO, 1017
EMMATTY, D. A., 375
ENGELHARD, ARTHUR W., 679; 851
EPLEE, ROBERT E., (861)
EVANS, G., (977)

FARLEY, J. D., 654, 673
FARMER, LOWELL J., 1136
FASSULIOTIS, GEORGE, 666
FAY, FLORENCE E., (471)
FEDER, W. A., (47)
FEHRMANN, H., (691)
FELICIANO, ALBERTO, (774)
FENTON, ROBERT, (241)
FENWICK, H. W., (501)
FERNALD, E. K., (546)
FILER, T. H., Jr., (93), 460
FILSINGER, ELIZABETH C., (33)
FIRSTMAN, ARANGA, (1136)
FISHER, C. D., (312)
FLEISHER, ZIPORA, 431
FLEMING, A. A., 473
FLETCHER, JOHN T., (120)
FOLEY, D. C., 855
FORBES, IAN, Jr., (10)
FORER, LYLE, (495)
FORSTER, M. J., (469), (938)
FRASER, L. R., (998)
FREDERICK, R. J., 223
FREDERIKSEN, R. A., 202
FRENCH, D. W., (682)
FRENCH, F. L., (291)
FRIDLUND, PAUL R., 738
FRIEND, DAVID, 885
FRY, W. E., (1085)

FUCIKOVSKY, L., 231
FULTON, N. D., 307, (879)
FUTRELL, M. C., 954

GABRIELSON, R. L., 234
GALLEGOS B., CESAR C., (203)
GARDINER, KATHLEEN D., 175
GARREN, KENNETH H., 419, (530)
GAY, J. DAN, 384
GERALDSON, C. M., (26)
GERDTS, MARVIN H., (69), (804)
GIBSON, KENNETH E., 85
GILL, C. C., 797
GILL, D. L., 731
GILL, H. S., 607
GILMER, R. M., (913)
GINN, ROBERT F., (1073)
GOLDEN, A. M., (114), (115)
GOODING, GUY V., Jr., 1013
GOTO, MASAO, 404
GOULD, C. J., (77), 425, 428
GOYAL, J. P., (752)
GRACEN, V. E., 469, 938
GRAHAM, DAVID A., 490
GRAVES, A. A., 810
GRAVES, C. R., (115)
GRAY, ELMER, 337
GRAY, F. A., (621)
GRAY, G. G., (959)
GRIFFIN, G. D., 1069
GRIMM, GORDON R., 929
GROGAN, C. O., 400, (469), 938
GROGAN, R. G., 3
GROSKOPP, M. D., (550)
GROSSMANN, F., (691)
GUDAUSKAS, ROBERT T., 947
GUILFORD, R. K., (234)
GUTHRIE, JAMES W., (54)

HAGEDORN, D. J., (127), 249, (408), 533, (735)
HALL, B. J., (841)
HALL, D. H., (757)
HAMMETT, K. R. W., 916
HANNA, M. K., (399)
HANSON, E. W., (127)
HANSON, JAMES B., 761
HARDER, D. E., 802
HARMAN, G. E., 639
HARMON, D. L., (871)
HARRIS, H. B., 312
HART, W. H., 89
HARTILL, WILLIAM F. T., 889
HARTWIG, E. E., (1043)
HARVEY, O. A., (449)
HAWKINS, ARTHUR, (968)
HEARN, C. J., 241
HEBERT, T. T., (956)
HEIT, C. E., (639)
HEWITT, W. B., 860
HIBBEN, CRAIG R., 475

HILDEBRANDT, A. C., (145), (1017)
HILDRETH, W. R., (1006)
HILTY, J. W., (115), 195
HITCHCOCK, S. W., (645)
HOCK, W. K., 58, (271), 971
HOFFMAN, J. A., 1132
HOLLAND, A. H., (449)
HOLMES, J., 643, 684
HOOPER, G. R., 824, 1108
HOPKINS, DONALD L., 610
HORNER, C. E., 814
HOWARD, C. M., 139, 142
HOWELL, W. E., 103, 676
HUDLER, GEORGE W., (1040)
HUNT, R. S., 659
HURLIMANN, JOSEPH H., (238), (586)

ISOM, W. H., (433)

JACOBSEN, B. J., 934
JAHN, OTTO L., (149)
JARES, T. W., (1073)
JAYNES, RICHARD A., 199
JENSEN, A. S., (390)
JOHN, C. A., (375)
JOHNSON, LEANDER F., 1126
JONES, A. C., 314
JONES, A. L., 961
JONES, B. L., 794
JONES, J. P., 1053
JONES, JOHN PAUL, 26
JONES, R. R., 1106
JONES, THOMAS W., (108)
JOSEPHSON, L. M., 115, (195)
JOSHI, L. M., (544), 1026
JUDD, R. W., Jr., 374, 1120
JUDGE, F. D., (457)
JUTTNER, A. S., (700)

KAIS, A. G., 309
KAISER, WALTER J., 244
KALOOSTIAN, G. H., (295)
KAMINSKI, W., (445)
KANTZES, JAMES G., (755), (1097)
KAPPELMAN, A. J., Jr., 896
KAVANAGH, T., (175)
KERNKAMP, M. F., (695)
KESSLER, KENNETH J., Jr., 788
KHOSLA, H. K., (95)
KILPATRICK, R. A., 871
KIMBLE, K. A., (3)
KINCER, H. C., (115)
KING, T. H., (498)
KINGSLAND, GRAYDON C., 118, 1122
KLEIN, M., (223)
KLINE, D. M., (325), 858
KNAPHUS, GEORGE, (855)
KNAUSS, J. F., 614
KNIGHTEN, JOHN L., 281
KOIKE, H., 766
KOMMENDAHL, THOR, 371, 726
KRAFT, J. M., 1056
KRAUSE, R. A., (757)
KRUSBERG, L. R., 276, (280)
KUHN, C. W., 453

LACEFIELD, GARRY D., (337)
LACY, M. L., (824)
LAEMMLEN, F. F., 743
LAI, MINGTAN, 1030
LAMBERTI, F., 111
LANDIS, Wm. RONALD, 634
LANG, D. S., (371), (726)
LAUGHLIN, CHARLES W., (167), (399)
LAUTZ, WILLIAM, (761)
LEACH, S. S., 723
LEAR, BERT, 1055
LEEPER, P. W., (794)
LEONARD, K. J., 759
LEWIS, J. A., (352), 440
LITTRELL, R. H., 951
LOEBENSTEIN, G., (431)
LOEGERING, W. Q., 718, (871)
LONG, J. K., (998)
LONG, P. G., 50
LORBEER, J. W., (394), (565), (1000)
LOWE, JAMES A., (337)
LUCAS, L. T., (3), 582
LUEPSCHEN, N. S., (314)
LUKENS, R. J., 645

MAAS, JOHN L., 72, 883, 1009
MADDEN, G., (72)
MAGGENTI, A. R., (89)
MAGIE, R. O., 82, 340
MAHMOOD, TARIQ, 464, 1130, 1132
MAI, WILLIAM F., (617)
MAJEED, M. G., (192)
MALEK, R. B., (927), (974)
MANION, PAUL D., 662, (1085)
MANNING, W. J., 47, 687
MARAMOROSCH, K., (223)
MARINOS, M. G., 682
MARSH, PAUL B., (33), (510), (714)
MARTIN, LLOYD W., 220
MARTIN, W. J., 523
MASSIE, LOWRY B., (495)
MATHUR, S. B., (349)
MATUSKEWSKI, MARK, (591)
MAXWELL, A. H., (281)
MAXWELL, DOUGLAS P., (361), 920
MAXWELL, JAMES D., (158)
McCAIN, ARTHUR H., 347
McCALLU, D. K., (399)
McCOY, R. E., 832
McCRACKEN, F. I., 93, (460)
McDONALD, R. E., 183
McDONNELL, P. F., 771
McELROY, F. D., 266
McFEELEY, J. C., 1064
McINTOSH, D. L., 213
McMEEKIN, DOROTHY, 877

MEISTER, CHARLES W., 861
MELUGIN, STELLA A., (322)
MELVIN, C. F., (149)
MERRILL, W., (1085)
MESNER, WILLIAM, (257)
MICHEL, L. J., 907
MILBRATH, G. M., 783, (785)
MILDENHALL, J. P., 536
MILLECAN, ARTHUR A., (555)
MILLER, H. N., 357, 587, (851)
MILLER, J. W., 38, 261
MILLER, P. M., (645), (968)
MILLER, PAUL ROBERT, 555
MILLER, V. L., 77, (425), (428)
MILLIKAN, D. F., (445), (828), (932)
MINK, G. I., (103), (676)
MINTON, N. A., 115
MITCHELL, J. E., (536), (1032)
MOHAN, S. K., (593)
MOHN, C. A., (460)
MOLLER, W. J., 1006
MORENO, M., (231)
MORGAN, OMAR D., 755, 1087
MORIN, CARL, (1030)
MORRISON, R. H., 498
MORSE, FRANK S., (648)
MORTENSEN, J. A., (610)
MOSSAHEBI, G. HOSSEIN, (244)
MOZINGO, R. W., (530)
MUKHOPADHYAY, A. N., 630
MULLER, IRMGARD, (874)
MUSE, R. R., 333
MUSSELL, HARRY, 471
MYERS, R., (1108)
MYSZKA, T., (445)

NAIK, RENUPAD, (96)
NATOUR, R. M., 192
NATTI, J. J., 457, 483
NAYAR, R., 170
NDUBIZU, THOMAS O. C., 791
NEELY, DAN, 37; 898
NEJA, RUDY, (860)
NELSON, J., (449), (841)
NELSON, JOHN W., 651
NELSON, R. R., 99, 325, 401, 495
NEMA, K. G., 95
NEMEC, S., 573
NEWBURG, W., (297)
NEWHALL, A. G., 216
NEWTON, O. H., (61)
NICHOLLS, THOMAS H., 1040, (1116)
NICHOLS, LESTER P., 509, (1085)
NICHOLSON, J. F., 911, 959
NUQUE, F. L., (17), (22)
NYVALL, ROBERT, (57)

O'BANNON, J. H., 154
OBRERO, F. P., 521
O'CALLAGHAN, T. F., (175)
OITTO, W. A., (379)
OKHOVAT, MAHMOUD, (244)
ONESIROSAN, P. T., 820
ORELLANA, R. G., 1124
ORILLO, F. T., (216)
ORR, C. C., 61
OSHIMA, N., 609, (613)
OU, S. H., 17, 22
OVERMAN, A. J., (26)
OYEKAN, P. O., 1032

PADDICK, R. G., 291
PAPAVIZAS, G. C., 352, (440)
PAPIA, P. M., (47), (687)
PARMETER, J. R., Jr., (659)
PARRIS, G. K., 778
PARTRIDGE, A. D., 1064
PATEL, P. N., 365
PATEL, V. C., 628
PATHAK, K. D., 544
PATHAK, V. N., 752
PAULUS, A. O., 449
PAVGI, M. S., (846)
PERKINS, I., (47)
PETERSEN, DONALD H., (509)
PETERSON, J. L., (374), (1120)
PETERSON, ROGER S., 163
PHILIS, J., 707
PHILLIPS, D. J., 837
PHIPPS, PATRICK M., (201), (467)
PINCKARD, J. A., (1111)
PITRE, H. N., (628)
PORTER, D. MORRIS, (419), 530
POWELL, DWIGHT, 96
POWERS, H. R., Jr., 623
PRATT, R. G., (536)
PRESCOTT, J. M., 695
PREY, ALLEN J., 648
PRICE, T. V., 67
PRITCHARD, A. J., (596)
PUCHALA, Z., (932)
PURCIFULL, D. E., 1061
PURUSHOTHAMAN, D., 107

RAABE, ROBERT D., 93, 238, 586
RADEWALD, J. D., 433, 841
RAINA, G. L., 283
RAMOS, D. E., (1006)
RAND, R. E., (249), (533)
RANDALL, W. K., (460)
RANNEY, C. D., 285
RAO, Y. P., 593
RATHAIAH, Y., 846
RAWLINS, T. E., (569)
REDDY, P. RANGA, (593)
REXRODE, CHARLES O., 108, 1106
REYES, L., (202)
RHOADES, H. L., 412
RIBEIRO, O. K., 599
RICH, AVERY E., 976
RICH, SAUL, (774)
RIEDEL, R. M., (673)

RIESSELMAN, J. H., (580)
RIFFLE, J. W., (823)
ROBERTS, D. A., 390
ROCHOW, W. F., 874
ROELFS, A. P., 986, (990), 992
ROSS, ELDON W., 741
ROTHMAN, P. G., (187), (992)
ROWELL, J. B., 990

SAAD, A. T., (1082)
SAAD, S. M., 735
SAETTLER, A. W., 703
SANTAMARIA, P. A., 349
SASIKALA, M., 185
SAYRE, K. D., (469)
SCHEIFELE, G. L., 302
SCHENCK, N. C., (486), 949
SCHIEBER, EUGENIO, 207, 209
SCHNATHORST, W. C., 780, 977
SCHOEN, JAMES F., 130
SCHOENHOLZ, JAMES, (591)
SCHREIBER, L. R., (58), (971)
SCHUSTER, M. L., (203), 505, (577), (580)
SCHWARZ, R. E., 747
SCOTT, D. H., (1009)
SCOTT, D. J., 966
SCOTT, GENE E., (954)
SCOTT, L. E., (1097)
SEAY, TIMOTHY, (697)
SELF, R. L., (424), (697)
SENECA, E. D., (582)
SETLIFF, EDSON C., 257
SHANMUGANATHAN, N., 1021
SHARMA, HARISH C., (752)
SHARMA, R. C., (544)
SHARP, E. L., (834)
SHEKHAWAT, G. S., (365)
SHIBUYA, F., (433), (449), (841)
SHIFRISS, CHEN, 604
SHOEMAKER, P. B., 565
SIBBETT, G. S., (780)
SIEMER, S. R., 297
SIGNORET, P. A., 1090
SILVA, J. P., (17), (22)
SIMONS, M. D., (907)
SIMPSON, MARION E., 33, 510, 714
SIMPSON, W. R., 501
SINCLAIR, J. B., (911), (959)
SINCLAIR, W. A., 1085
SINGH, A. B., 526
SKILLING, DARROLL D., 1116
SMITH, HERBERT L., 13
SMITH, J. DREW, 63
SMITH, R. R., (920)
SMITH, ROBERT L., 1099
SMITH, S. H., (295)
SMITH, W. L., Jr., 228
SMOOT, JOHN J., 149
SNEL, MARTEN, 120
SNOW, G. A., (764)
SNYDER, W. C., (837)
SOLEL, Z., 408
SOUTHARDS, CARROLL J., 41
SOWELL, GROVER, Jr., (312)
STALEY, J. M., (764)
STAMBAUGH, W. J., (700)
STATLER, GLEN D., 172
STEELE, A. E., 1101
STEEPY, T. L., 751
STEVENSON, W. R., 408
STEWART, D. M., 187
STIPES, R. JAY, 201, 467
STOVER, R. H., 437
STRANDBERG, J. O., 142
SUBIRATS, F. J., 424, 697
SUIT, F. R., 923
SUMNER, DONALD R., 540, (951)
SUNDARAM, N. V., (87)
SUTABUTRA, T., 328

TAINTER, F.H., (1036)
TANVEER, M., 1082
TATE, C. J,. (1077)
TAYLOR, P. A., 361
TEAKLE, D. S., 596
TEEM, DAVID H., (947)
TEMPLETON, G. E., (226)
THAKUR, R. P., (630)
THOMAS, C. A., 1046
THOMPSON, D. L., 158, 956
THOMPSON, J., (1101)
THYR, B. D., 336
TOMERLIN, A. T., (154)
TOMLINSON, HARLEY, 774
TO, WAI-NAM, (947)
TONTYAPORN, S., 226
TOWERS, BARRY, 591
TOWNSEND, ALDEN M., 980
TRAYLOR, JACK A., 294
TRUE, R. P., (1106)
TRUJILLO, E. E., (521)
TUITE, JOHN, 387
TURNER, P. L., (291)

ULLSTRUP, A. J., (202), 563, 1046
UYEMOTO, J. K., 913

VALENTINE, FREDERICK A., (662)
Van ARSDEL, E. P., 1073
van der ZWET, T., (72), 379
van SCHAIK, P. H., (530)
Van VUUREN, S. P., (747)
VARGAS, J. M., Jr., 167, (399), 1118
VAUGHAN, EDWARD K., (57), (297)
VEST, G., (824)
VILLENA, WILLY L., (158)

WAGNON, H. KEITH, (294)
WALKER, JERRY T., (475), 893
WALKEY, D. G. A., 268
WALLIN, JACK R., 479
WALTON, G. S., (645)

WARD, H. N., (1064)
WARMKE, H. E., 486
WARREN, H. L., (1077)
WARREN, T. B., (582)
WATERWORTH, H. E., 983
WATSON, A. G., (837)
WATSON, D. R. W., 517
WATTERSON, J. C., 816
WEAVER, D. J., 1094
WEBB, RAYMON E., 132
WEBER, G. F., (390)
WEBSTER, R. K., 757
WEHLBURG, C., 447
WEHUNT, E. J., (415)
WEIGLE, C. G., (1030)
WEIHING, J. L., 580
WEINHOLD, A. R., 837
WEISS, M. J., 823
WELLS, HOMER D., 10
WELLS, JOHN M., 69, 804
WESTDAL, P. H., (802)
WHATLEY, THOMAS L., (947)
WHEATLEY, G., (1101)
WHEELER, HARRY, 667
WHITE, DONALD G., 382
WHITNEY, E. D., 30
WICK, C. M., (757)
WILCOX, J. R., 776
WILLIAMS, A. S., (667)
WILLIAMS, CURTIS, (1043)
WILLIAMS, EUNICE, (913)
WILLIAMS, HAROLD E., (294), 295
WILLIAMS, P. H., (536), (816), (934)
WILLIAMS, R. A., 1036
WILSON, K. I., (185)
WILSON, V. E., 730
WITCHALLS, J. T., 45
WITCHER, WESLEY, 158, (810)
WOODHOUSE, W. W., Jr., (582)
WUEST, P. J., (684)

YARWOOD, C. E., 342
YOUNG, L. D., (667)

ZAGAJA, S. W., 445
ZEHR, ELDON I., (158)
ZENTMYER, GEORGE A., (207)
ZETTLER, F. W., (506)
ZIMMER, D. E., 11
ZITTER, T. A., (1061)

SUBJECT INDEX

Acacia melanoxylon: root rot disease, caused by Poria vincta var. cinerea, in East Africa for the first time, 257

Adjuvants: 23 different combinations of adjuvants to benomyl did not increase its uptake by American elm seedlings in planting medium, 971

Aerial spray: aerial sprays of phytoactin and semicarbazone ineffective in control of Hypoxylon canker on aspen in Minnesota, 650

Aesculus spp.: additional species and subspecies found susceptible to leaf blotch, caused by Guignardia aesculi (= Phyllosticta sphaeropsoidea), 37

Africa: 209; 257; 672; 747; 820

Agaricus bisporus (mushroom): benomyl and thiabendazole for control of mushroom diseases, 120; effect of time of inoculation on incidence and control of Verticillium disease, 643

Agrobacterium tumefaciens: 145

Air pollution: extent of oxidant air pollution damage to some pines and other conifers in California, 555

Alabama: 424; 697; 947

Alfalfa: seedlings inoculated with Ditylenchus dipsaci grown in presence of dichlorvos pest strip showed reduced nematode numbers, 621

Allium spp. (see Onion)

Alternaria sp.: 714; 791; 848

--- brassicicola: 454

--- solani: 807

--- tenuis: 107; 297; 639; 776; 864

Angular leaf spot: of strawberry, caused by Xanthomonas fragariae, first report from Florida, 142

Announcement: Monograph No. 6, "Fusarium wilt of tomato," by J. C. Walker, now available through the American Phytopathological Society, 653

Anther callus: production of virus-symptomless geranium plants from anther callus, 1017

Anthracnose: on Dolichos biflorus, in Georgia, 10; of sycamore, severe defoliation and twig blight in epidemic proportion in the Mississippi-Yazoo Delta, 93; of strawberry, benomyl + pinolene gave control in Florida if applied twice weekly, 139; of tomato, inheritance of resistance appears to be partially dominant with several genes involved, 253; of corn in Maryland, caused by Colletotrichum graminicola, 755; stem anthracnose of cowpeas in Nigeria, 820; northern anthracnose of red clover, development of a population of red clover with a higher degree of resistance to Kabatiella caulivora than exists in present cultivars, 920;

of black currant, controlled in Poland with benomyl, 932
Anthriscus sylvestris (cow parsley): one of natural weed sources of western celery mosaic virus in England, 268
Antibiotics:(see also Chemicals and Fungicides): phytoactin and semicarbazone failed to control Hypoxylon canker on aspen in Minnesota when applied aerially, 650; streptomycin sulfate gave fairly good control of bacterial leaf spot on Ixora coccinea, in Florida, 261
---: tetracycline-hydrochloride, a bioassay described for determination of activity of this antibiotic in plant extract, 223; suppression of Pierce's disease of grape effected by tetracycline antiobiotics in Florida, 610; injection of tetracycline-hydrochloride solution into trunks of greened sweet orange trees reduced greening, 747
Antirrhinum majus (snapdragon): lesion nematodes controlled in planting soil in greenhouse with aldicarb and BAY 68138, 276; fungicide control of Phytophthora in seedlings with ETMT, 587; Alternaria tenuis and Phyllosticta antirrhini deeply borne within seed are pathogenic to seed and seedlings, 639
Aphanomyces cochlioides: 440
Aphid-borne viruses: of sugarbeet, their incidence and distribution in Washington, 103
Apple: apple twig canker in Illinois, caused by Monochaetia mali, 96; fruit rot, from Botryosphaeria ribis and Physalospora obtusa under orchard conditions, major source of inoculum is pycnidiospores from pycnidia on dead twigs, 122; crown rot, dilution plates show Dexon, Difolatan and maneb to be most suitable of fungicides tested for soil injection to prevent crown rot infection from Phytophthora cactorum, 213; core rot, caused by Trichothecium roseum found in apple orchards in India; previously considered a postharvest disease, 283; Nectria canker, field resistance to Nectria galligena observed in Beauty of Rept, Boiken, Close and Wealthy varieties, 445; nematodes, Pratylenchus penetrans controlled in New York orchards with D-1410, 617
---: crab: the imperfect stage of apple scab fungus overwintered on varieties of flowering crab apple in Pennsylvania, 509; inoculation of Virginia crab apple with pear latent virus incited symptoms of virus, 983
---: powdery mildew: caused by Podosphaera leucotricha, effectiveness of fungicides became greater with seasonal increase in temperature, 263; benomyl and triarimol compared favorably with Karathane in control of powdery mildew in Washington trials (2 years), 514
---: postharvest decay: from Nectria, controlled with 2-aminobutane dip, 771; postharvest fungicide treatment of thiabendazole on mechanically harvested 'McIntosh' apple very effective, 791
---: scab: control of apple scab, bitter rot and sawfly in Poland with some newer fungicides, 828; conidia of Venturia inaequalis frozen and stored as long as a year retained infectivity, 976
Application dates: comparison of application dates for control of Typhula blight of creeping bentgrass with fungicides, 1118
Apricot: Eutypa armeniacae, cause of apricot limb dieback, attacks Ceanothus at pruning wounds, in California, 1006; stem pitting of peach and apricot found recently in Greece, 1049
Arabidopsis thaliana (mouse-ear cress): common weed in alfalfa fields found to be host of Sclerotinia sp., 1087
Arachis hypogaea (see Peanut)
Areca cathecu (arecanut): yellow leaf disease, mycoplasma colonies found associated with infected leaves of arecanut, in India, 170
Arkansas: 226; 307; 411; 1036
Armillarea mellea: 823; 1099
Ascochyta sp.: 1064
Aspen (See Populus)
Aspergillus spp.: 804
--- flavus: 33; 510; 714
--- niger: 714
Australia: 67; 291; 596; 977; 998
Azalea: benomyl as a cutting soak or cutting drench controlled Cylindrocladium rot, 679

Bacteria: identification of different bean blight bacteria by bean seedling injection, 703
Bacterial diseases: angular leaf spot of strawberry in Florida, caused by Xanthomonas fragariae, 142; of garden pea, in New Zealand, two Pseudomonas isolates were isolated from affected plants, 517
---: bacterial blight: of barley reported to occur in India for the first time, 365; common blight of bean, new highly virulent strains of Xanthomonas phaseoli obtained from dry bean seed from Colombia, 505; common blight and fuscous blight of bean, disease reaction of bean varieties tested by two inoculation methods, 577; caused by Xanthomonas phaseoli, copper bactericides gave appreciable control even after beginning of secondary spread, 609; separation of bean blight bacteria by

seedling injection, 703; oleander blight of oleander in California, 1030; of rice, over 8000 rice varieties screened by artificial inoculation for resistance to Xanthomonas oryzae isolates in the Philippines, 17; Indian isolates of Xanthomonas oryzae from rice differed greatly in degree of virulence, 593
---: bacterial canker: of Prunus spp. orchard trees in Oregon, 421; of sweet cherry in Michigan, 961; of tomato, Solanum triflorum found to be natural weed host in Wyoming, 336; recovery of Corynebacterium michiganense from overwintered tomato stems by the excised-petiole inoculation method, 654
---: bacterial brown spot of bean, control with cupric hydroxide (K 101), 735
---: bacterial leaf spot of Ixora coccinea, caused by Xanthomonas maculifoliigardeniae, in Florida, 261
---: bacterial leaf streak of rice, an inoculation technique for testing virulence of bacterium or resistance of host plant, 404
---: bacterial leaf stripe of Strelitzia reginae, in Florida, 447
---: bacterial soft rot and black spot, of bok choy (Brassica campestris), main causes of market losses, 1037
---: bacterial spot of peach, previously undescribed symptom, a gray colored leaf spot is caused by Xanthomonas pruni, 774
---: bacterial wilt: of cucurbits, 9 of 10 resistant species were from the genus Cucurbita, 816; of tomato, in Hawaii, controlled by grafting to resistant rootstocks, 521
---: crown gall, retardation of crown gall enlargement of tomato after bacteriophage treatment, 145
---: halo blight; of bean, Pseudomonas phaseolicola found in only three fields of same variety and source; drastic decline in incidence in Idaho attributed to an effective program of inspection, regulated importation of seed and seed application of 5% slurry of streptomycin sulfate, 54; inheritance and linkage of halo blight systemic chlorosis and leaf watersoaked reaction in bean variety crosses, 203; of timothy, caused by a Pseudomonas sp. in Wisconsin, 361
Bacteriophage treatment: absorption and translocation of PB2-A bacteriophage of Agrobacterium tumefaciens by tomato plants and retardation of stem gall development after its absorption, 145
Banana: black leaf streak, caused by Mycosphaerella sp. controlled with benomyl, in Samoa, 50; banana leaf spot (Sigatoka and black leaf streak), two distinct diseases caused by different pathogenic forms of M. musicola from Central America and the Philippines, respectively; contrasting features of control are outlined, 437; control of root-knot and spiral nematodes on banana in Cyprus by fumigation with DBCP, 707
Barbarea vulgaris (yellow rocket): host of Sclerotinia sp. in alfalfa fields, 1087
Barberry (Berberis thunbergii atropurpurea): increase in Verticillium infection after heat treatment of soil, 996
Bare-root dips: nematicidal dips effective in control of soybean cyst nematode, 13
Bark and wood pitting disease: of grape, observed for first time in California in 1971 and is believed to be caused by a strain of grape fanleaf virus, 860
Barleria cristata: leaf blight is caused by Alternaria tenuis in India, 107
Barley: seed treatment before storage, hot water and anaerobic treatments reduced seedling emergence, 118; bacterial blight of barley caused by Xanthomonas translucens f. sp. hordei, in India, 365; control of covered smut of barley with carboxin seed treatments, 544; resistance to Helminthosporium stripe in winter barley cultivars, 858; leaf rust, caused by Puccinia hordei, sensitivity to external environment and physiologic specialization, 1026; several fungicides tested for control of loose smut of barley and barley leaf stripe, 1122
---: virus: barley yellow dwarf: prevalence of aphids and BYDV in Manitoba in 1970, 797; a fifth variant of barley yellow dwarf virus has been found in New York, 874
Bean: one of four selective hosts utilized to identify strains of alfalfa mosaic virus, 127; new bean disease found in Caribbean area, caused by a species of Entyloma, 207; field control of white and gray molds of beans in Washington with benzimidazoles, DCNA, and cyanamide, 234; black root rot, caused by Thielaviopsis basicola, controlled with soil amendments and fungicides in the greenhouse, 352; defoliation, caused by rain damage at primary leaf stage, in New York, 457; seed and soil treatments for control of bean root rots, mainly Fusarium solani f. phaseoli, 483; bacterial brown spot, copper fungicide as foliage spray gave good control, 735; Pythium ultimum as a root pathogen of bean and peas in Washington, 1056
---: blight: two Colombian isolates of Xanthomonas phaseoli, cause of common blight, are apparently new strains of pathogen, 505; two inoculation methods used to compare the disease reaction

of Phaseolus vulgaris and P. acutifolius to common blight and fuscous blight bacteria, 577; copper sprays for secondary spread of common blight, 609; a seedling injection method to identify different bean bacteria, 703
---: halo blight: found in only three fields in Idaho, the control program in force has drastically reduced infection, 54; inheritance and linkage of the halo blight systemic chlorosis and leaf watersoaked reaction in Phaseolus vulgaris variety crosses, 203
Bedding plants: Pythium and Phytophthora infection controlled in bedding plants of petunia, snapdragon, Calendula, Dianthus, Verbena, and Vinca with ETMT fungicide as a soil drench or soil mix, 587
Beech: beech bark disease, the presence of beech scale (Cryptococcus fagi) infestation in northeastern Pennsylvania beech stands indicates the potential for the Nectria component of the disease to cause severe loss in the near future, 591
Belonolaimus longicaudatus: 412
Bioassay: fungitoxic activity of benomyl in American elm shoots and twigs against Dutch elm disease, 179; antibiotic tetracycline-hydrochloride detected in measurable concentrations in plant extracts from China aster up to 10 days after treatment, 223
Biological control: of forest root pathogens by stump treatment with Trichoderma sp. in chain oil of chain saw, 659
Birch (Betula alleghaniensis): stem abnormalities of yellow birch seedlings in Michigan and Wisconsin, 788
Bitter rot: of apple, controlled with some newer fungicides, in Poland, 828
Black leaf streak: of banana, controlled in Samoa with benomyl in oil-water emulsion, 50; in Central America and the Philippines, 437
Black pod: of cacao, caused by Phytophthora palmivora, caused serious losses in all areas of the Solomon Islands, 885
Black root rot disease: of strawberry, caused by Rhizoctonia fragariae, evidence indicates environmental and/or nutritional conditions influence pathogenicity, 599
Black spot disease: of bok choy, caused by Erwinia carotovora, 1037
Blight: Ascochyta, of cotton, estimate of loss by State in 1970, 504
---: bacterial: of rice, virulent isolates of Xanthomonas oryzae from Philippines tested in 8700 rice varieties, none immune but some showed resistance, 17; of barley, "Xanthomonas streak" or bacterial blight reported from India for first time, was also pathogenic to rice, 365; estimated loss of cotton from Xanthomonas malvacearum in 1970, 504; of garden pea, in New Zealand, 517; of oleander, found in California, 1030
---: ---: of bean: new virulent strains of Xanthomonas phaseoli are present in Colombia, 505; leaf water-soaking was more effective than stem inoculation to compare the disease reactions of bean varieties to common blight or fuscous blight bacteria, 577; 8% metallic copper from copper ammonium carbonate gave good control of common blight in Colorado and can be used by aircraft, 609
---: early blight, of tomato, caused by Alternaria solani, field evaluation of 30 tomato varieties or lines for resistance, 807
---: Fusarium blight: of 'Merion' Kentucky bluegrass controlled with benomyl, 167; benomyl and thiophanate were effective on naturally infected 'Merion' Kentucky bluegrass, 333
---: halo: of bean, only three fields in Idaho found infested with halo blight, from seed grown in 1967, 54; bean variety crosses studied to determine the controlling factors in their inheritance of systemic chlorosis and watersoaked leaf reaction to Pseudomonas phaseolicola, 203; of timothy, in Wisconsin, 361
---: leaf: of Barleria cristata in India, caused by Alternaria tenuis, 107
---: ---: of onion, resistance and susceptibility of Allium species, commercial varieties and interspecific crosses to Botryotinia (Botrytis) squamosa), 394; controlled with with maneb with 7-day schedule, 565
---: ---: southern corn leaf blight (SCLB): survival and overwintering properties of isolates of Helminthosporium maydis, 99; reductions in yield because of blight, 115; distribution of Race T in the United States in 1969 and 1970, 302; naturally infected kernels, 371; pathogen found to be carried on popcorn seed from planting infected with SCLB, 382; seed infection in the seed samples from the 1970 crop in Indiana, 387; spread in Hawaii indicates infected seed lots as cause, 392; selection of plants resistant to Race T of H. maydis by rapid testing method, 469; effect of cytoplasm on resistance or susceptibility 473; the frequency and mating types of H. maydis by race and virulence in 1970 in Northeastern U. S., 495; overwintering of Race T, 563; use of T-toxin as indicator of resistance to SCLB, 667; reported from Nigeria in 1970, 672; Race T on seed produced in Iowa in 1968, 855; overwintering studies in five south-

ern States in 1970, 947; 949; 951; 954; 956; gift packs of popcorn from the U.S. imported to New Zealand found to be infected with H. maydis, 966
---; ---: yellow leaf blight: evaluation of various corn inbreds in normal and different male-sterile cytoplasms to Phyllosticta sp., 401; isolates C, U, and OR of yellow leaf blight compared as to pathogenicity and other characters; identity as to species is uncertain, 1064
---: Marasmius blight, of American beachgrass, along Outer Banks of North Carolina, 582
---: needle: of Scotch pine, brown spot, caused by Scirrhia acicola, has seriously damaged Scotch pine Christmas tree plantations in Wisconsin, 648; needle blight of southern pines is presumed to have been caused by Lophodermella, Ploioderma lethale and infrequently by Ploioderma hedgcockii, 764; of Austrain pine, in Minnesota caused by Dothistroma pini, 1040
---: petal blight, of camellia, caused by Pestalotia sp., in Mobile County, Alabama, 697
---: pod and stem, of soybean, marked differences in blight were associated with plant density, 776
---: Rhizoctonia blight, of swordfern, controlled effectively with benomyl, thiabendazole or chlorothalonil, 614
---: of Spanish moss, caused by a form of Fusarium solani, has caused extensive blighting of Spanish moss from Bradenton, Florida to Jacksonville and also occurs in southeastern Georgia, South Carolina and Mississippi, 390
---: Typhula blight, of bentgrass, controlled with chloroneb and a granular formulation of mercurous chloride and mercuric chloride, in Michigan, 1118
---: white thread blight and horse hair blight, of cacao, caused by Marasmius spp., 885
Blueberry: resistance to mummy berry disease, caused by Monilinia vaccinii-corymbosi, in named varieties and selections, 651
Bok choy (Chinese chard = Brassica campestris): transit and market losses caused mainly by Pseudomonas marginalis and Erwinia carotovora, 1037
Botryodiplodia theobromae: 152
Botryosphaeria ribis: 122
Botryotinia squamosa (Botrytis): 394
Botrytis cinerea: 135; 234; 791; 864; 883; 1009
--- squamosa: 565
Brassica oleracea (see Crucifers)
Brazil: 209
Bremia lactucae: 794
British Columbia: 266
Broccoli (see also Crucifers): sugarbeet cyst nematode, controlled with 1,3-D and DD applied at three depths, 1101
Brown root: of muck-grown carrots, caused by Pythium spp., 536
Brown rot: of peach, hot water dips or hot suspensions of DCNA or benomyl gave better control than unheated water or unheated chemical suspensions, 228
Bulb soak treatments: analytical reactions for a field test for thiabendazole concentration in treating tanks, 77
Bunt: common, control of seedborne and soilborne Tilletia caries with systemic fungicides benomyl and carboxin, 1132
---: dwarf bunt: of wheat, carboxin and formulations of thiabendazole evaluated as seed treatments, 613; some control of dwarf bunt of wheat was obtained with benomyl and carboxin applied as sprays to plants, but not as seed treatments, 1132
---: hill bunt of wheat, caused by Tilletia foetida and T. caries; benomyl seed treatment completely checked disease in India, 544

Cabbage (see Crucifers)
Cacao: incidence and importance of diseases of cacao in the British Solomon Islands Protectorate, 885
Calendula: control of Phytophthora in seedlings with fungicide ETMT, 587
California: 3; 30; 69; 89; 93; 294; 322; 433; 449; 490; 555; 607; 659; 757; 780; 804; 841; 860; 977; 1006; 1030; 1099; 1101
Calonectria sp.: 743
Cambial zone discoloration: caused by stem pitting, in peach and apricot, 1049
Camellia: new fungus petal blight identified as Pestalotia sp. found in Alabama, 697
Canker: of yellow birch seedlings in Wisconsin and Michigan, 788
---: bacterial: of Prunus, severity on orchard trees is related to the susceptibility of root or trunk stock, 421; of sweet cherry in Michigan, 961
---; ---: of tomato: a common weed (Solanum triflorum) found naturally infected with Corynebacterium michiganense, 336; in Ohio, C. michiganense recovered from overwintering stems of tomato by the vascular inoculation method, 654
---: Cytospora canker on peach, its seasonal development in Colorado with greatest enlargement during the spring, 314
---: Endothia: first report of inhibition of chestnut blight fungus in the field with a fungicide (benomyl), 199; cause of canker disease of pin oak in Hampton, Virginia, 467

---: Gloeosporium canker of Burford holly, reported for first time as causal agent on this host, 424
---: of Katsura-tree, Dothiorella ribis was isolated and caused wood discoloration when inoculated on wounded seedlings, 893
---: Monochaetia mali: caused apple twig canker in Illinois, 96; Monochaetia canker of Arizona cypress and redcedar in South Carolina, 810
---: Nectria canker of apple, varieties Beauty of Rept, Boiken, Close and Wealthy showed no infection under field conditions, 445
---: Phytophthora canker of cacao, in Solomon Islands, 885
---: Septoria canker of eastern cottonwood nursery stock, caused by Septoria musiva, in Mississippi, 460
---: stem, of tea plant, benomyl shows promise as a fungicide for control of tea stem canker, caused by Phomopsis theae, 1021
Cantaloup (see Cucurbits)
Capsella bursa-pastoris (shepherd's purse): weed host of Sclerotinia sp. in alfalfa fields, 1087
Carrot: nematode-induced abnormalities of carrot in southern Italy, 111; alfalfa mosaic virus found in carrots near Modesto, California, first report on this host, 322
Cauliflower (see Crucifers)
Ceanothus spp. (California wild lilac): Eutypa armeniacae (pathogen of apricot dieback) is cause of limb dieback in Ceanothus, 1006
Central America: 437
Centrospora acerina: 731
Ceratocystis adiposa: 7
--- fagacearum: 108; 281; 1106
--- fimbriata: 523
--- --- f. platani: 741
--- paradoxa: 7
--- ulmi: 158; 179; 898; 971; 980
Cercospora sp.: 923
--- beticola: 408; 449; 479; 1073
--- brassicicola: 540
--- carthami: 846
--- exitiosa: 733
--- kikuchii: 776
--- microsora: 733
--- omphakodes: 1120
Cercosporella herpotrichoides: 45
Cereal enation disease: of oats, in Kenya, 802
Ceylon: 1021
Chemicals (see also Antibiotics, Fungicides, and Nematicides): borax was effective in preventing colonization of Fomes annosus in ponderosa and Jeffrey pine stumps in California, 490
---: boron, excess boron responsible for tree collapse of citrus in Turkey, 1132
---: CO_2, latent Fusarium infections in gladiolus corms developed quickly into rotted tissue after 4 days in 50% CO_2 atmosphere, 340
---: chlorine, chlorine used with thiabendazole improved effectiveness as a postharvest fungicide for apples, 791
---: copper bactericides: used as protective sprays for secondary spread of common bean blight in Colordao, 609; tribasic copper sulfate + a coordination product of zinc ion and manganous ethylenebis [dithiocarbamate](copper-maneb) gave fairly good control of Xanthomonas maculifoliigardeniae on tropical shrub, Ixora coccinea, 261
---: ethephon ((2-chloroethyl)phosphonic acid), for gladiolus cormels, added to benomyl or thiabendazole in hot water yielded larger and heavier corms, 82
---: ethoxyquin and diphenylamine, storage scald inhibitors, used with thiabendazole proved compatible in postharvest treatment of 'McIntosh' apple, 791
---: herbicides: 2,4-dichlorophenoxyacetic acid (2,4-D), 4-amino-3,5,6-trichloropicolinic acid (picloram) + 2,4-D, and dimethylarsenic acid + arsenic (cacodylic acid) not effective in preventing mat formation or oak bark beetle attack in oak wilt trees, 1106
---: insecticides: systemic organophosphorus types, cynem and phorate, dimethoate, and oxime carbamate insecticides reduced incidence of tobacco yellow dwarf; hypothesis proposed that possibly the chemicals act directly on the causal organism as well as the vector, 291; D-1410 (S-methyl 1-(dimethylcarbamoyl)-N-[(methylcarbamoyl)-oxy] thioformimidate) a nematicide-insecticide on a 4-spray schedule to apple, peach, pear and cherry trees, reduced Pratylenchus penetrans in roots by more than 90% in New York State, 617; carbofuran (2,3-dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate) an insecticide-nematicide reduced stalk rot of corn by reduction of insect and nematode wounds, 634; dichlorvos (2,2-dichlorovinyl dimethyl phosphate), dichlorvos pest strips reduced stem nematode damage in seedling alfalfa, 621
---: nitrogen: NH_4OH and urea added to greenhouse soil reduced damping-off of sugarbeets, 440; liquid nitrogen storage of urediospores of Puccinia graminis f. sp. tritici without loss of viability has been successful over 10-year period, 871
Chemotherapeutants: carboxin and thiabendazole evaluated for control of dwarf bunt

of wheat by seed treatment, 613
Chenopodium spp.: Sphaerotheca lanestris and Erysiphe graminis transmitted a virus to Chenopodium quinoa which resembled tobacco mosaic virus, 342
Cherry (see also Prunus): Eton-Bug disease of sweet cherry, evidence suggests that the disease is caused by an undetermined virus, 344; control of Pratylenchus penetrans with foliar sprays of D-1410, 617; bacterial canker of sweet cherry reported for the first time from Michigan, 961
Chestnut (Castanea dentata): inhibition of Endothia parasitica by benomyl in field-grown American chestnut trees, first report of a fungicide that inhibits growth of Endothia in the field, 199
China aster (Callistephus chinensis): acquisition and retention of tetracycline-hydrochloride in leaves, 223
Chinese gooseberry (Actinidia chinensis): potential new crop, found to be infected by Armillaria mellea in California, 1099
Chokecherry (Prunus virginiana): eradication of infected chokecherry from around peach and nectarine orchards for control of X disease, a major peach disease in the Northern United States, 645
Christmas tree diseases: brown spot needle blight of Scotch pine, 648; needlecast of Scotch pine, 1116
Chrysanthemum: Pratylenchus penetrans controlled in heavy soil in greenhouse for at least 10 months with use of nematicides aldicarb and BAY 68138, 276; rooted cuttings sprayed or dipped in benomyl suspensions were immunized to infection by Verticillium albo-atrum and Rhizoctonia solani, 347; Pythium root rot, of potted plants controlled with ETMT, 851
Cicadulina mbila: 802
Circulifer opacipennis: 85
--- tenellus: 85
Citrus: nematodes, Meloidogyne javanica found on sweet orange, sour orange, Troyer citrange and Cleopatra mandarin rootstocks in California, 607
---: Phytophthora, fumigation of Phytophthora in sandy soil by surface application of methyl bromide and methyl bromide-chloropicrin before planting citrus, 929
---: pink pitting on grapefruit, caused by Cercospora sp. controlled with 1.3% oil, zineb or neutral copper, in Florida, 923
---: postharvest decay, washing oranges and tangerines and applying fungicides before degreening was more beneficial in reduction of postharvest decays than the standard procedure of washing and treating after degreening, 149
---: sour orange scab, on 'Page' citrus hybrid, benomyl spray prevented scab on leaves and fruits for periods up to 150 days, and proved better than copper spray, 241
---: tree collapse: in sweet orange and mandarin in Turkey, cause unknown, 464; boron toxicity is cause of tree collapse in Izmir region of Turkey, 1132
---: virus: decrease in fruit greening of sweet orange by trunk injection of tetracyclines, in South Africa, 747; exocortis virus in dwarfed citrus trees in Australia, 998
Cladosporium herbarum: 714
Claviceps gigantea: 231
Clover: isolate of Phytophthora parasitica from white clover inoculated on four soybean varieties may have limited usefulness for varietal verification, 130; development of red clover germ plasm resistant to northern anthracnose, caused by Kabatiella caulivora, 920; yellow sweet clover, red clover, and two white clovers found to be good hosts of lespedeza cyst nematode, 927
Cochliobolus carbonum (Helminthosporium sp.): 325
Code system: for cereal rusts, an infection-type coding system is proposed in which relative size of the sporulating area and relative total size of typical lesions are coded numerically and the nature of tissue damage associated with typical lesions is coded descriptively; uniformity of infection types from plant to plant is coded with a fourth code, 319
Coffee: rust, comparison of coffee rust in Brazil and Kenya, Africa, the pathogen, symptomatology, ecological aspects and control measures, 209; in the Philippines, Bordeaux and cuprous oxide sprays gave good control; care was taken to cover underside of leaf (stomatal location) where rust spores penetrate, 216
Cold pox: of butternut squash, in Florida, caused by low temperature, 350
Colletotrichum coccodes: 253
--- dematium f. sp. truncata: 10
--- fragariae: 139
--- gloeosporioides: 149
--- gossypii: 714; 879
--- graminicola: 755
--- higginsianum: 540
--- lindemuthianum: 820
Colocasia esculenta (dasheen and taro): virus infections were found in a large percentage of plants, in Puerto Rico, 506
Colorado: 314; 609
Concentric ring pattern: on seedling pear trees, no virus or pathogen known to cause the disorder, some evidence indicated for possible genetic inheritance, 379

Conifers: the current distribution and severity of oxidant air pollution injury to some conifers in California, 555
Core rot: of apple, caused by Trichothecium roseum, in preharvested apples in Kulu Valley, India, 283
Corn: high-lysine corn grown in 1971, because of demand for corn in normal cytoplasm, has shown extreme susceptibility to kernel and ear rots, 1046
---: anthracnose, heavy infection on corn in Maryland thought to be first observation on corn in Maryland, 755
---: Claviceps gigantea infection in Mexico on two lines of corn, fungus favored by mean annual temperatures of 13-15°C and over 1000 mm annual precipitation, 231
---: Diplodia stalk and ear rot, correlations between these two diseases indicate that they are generally independent in character, 158
---: downy mildew, resistance to Sclerospora sorghi found in corn inbred lines Tx441, Tx601, B46, H52, and Oh43, in field tests in Texas, 202
---: head smut, caused by Sphacelotheca reiliana, suppressed with carboxin, 501
---: kernel discoloration of 'Jubilee' sweet corn reported from western Washington, apparently a new disease not yet identified as to cause, 57
---: nematode density, crop host had greater influence on nematode density than fall tillage, 41
---: stalk, rot, carbofuran reduced stalk rot apparently by reduction in number of insect and nematode wounds, 634
---: southern corn leaf blight (SCLB): the overwintering and survival of isolates of Helminthosporium maydis, 99; yield reductions from SCLB, 115; reaction of corn inbreds with different cytoplasms to H. maydis, 195; distribution of Race T, 302; seedling wilt from kernels naturally infected with H. maydis, 371; high percentage of popcorn seed from planting severely affected with SCLB were infected with H. maydis, 382; infection of corn seed in Indiana, 387; spread in Hawaii, 392; resistance to H. maydis, Race T, 469; effect of cytoplasm on susceptibility to SCLB, 473; occurrence of Fusarium moniliforme and H. maydis on and in corn seed as related to T cytoplasm, 486; identification of 934 H. maydis isolates as to race frequency, distribution and virulence, 495; overwintering of Race T in the Midwest in 1970-71, 563; H. maydis toxin as indicator of resistance, 667; in Nigeria on widely grown cultivar NS-5, 672; detection of commercial samples produced in 1970 infected with H. maydis, 726; formation of sclerotia of H. maydis Race T on corn seed coat, 755; association of virulence and mating type among H. maydis isolates collected in 1970, 759; H. maydis Race T was found in kernels of maize produced in 1968 in three Iowa counties, 855; reactions of corn genotypes and cytoplasms to H. maydis toxin, 938; H. maydis overwintered and remained infective in five southern States in 1970, 947; 949; 951; 954; 956; seed lines and gift packs of popcorn imported to New Zealand found to be infected with H. maydis; an embargo placed on all imports of corn seed and raw corn products from the U.S., 966;
---: yellow leaf blight: reactions of corn inbreds in different cytoplasmic backgrounds to Phyllosticta sp., causal agent of yellow leaf blight, 401; three isolates were compared and their classification to genus is questioned, 1064
Correction: (Vol. 54, No. 2, Page 134), 464
Correction: (Vol. 54, No. 9, Page 745, 747), 96
Correction: (Vol. 55, No. 1, Page 58) 2nd line of Abstract should read "by soil and trunk injections did not reduce symptoms" instead of "by soil and trunk infections did not reduce symptoms"
Correction: (Vol. 55, No. 4, Pages 347, 348), 560
Correction: (Vol. 55, No. 5, Page 461), 656
Correction: (Vol. 55, No. 6, Pages 512, 513, and 521), 752
Correction: (Vol. 55, No. 7, Page 586), 944
Correction: (Vol. 55, No. 7, 643, 644, and 645) change "Verticillium malthausei" to "Verticillium malthousei"
Correction: (Vol. 55, No. 8, Pages 662, 663 664), 1040
Correction: (Vol. 55, No. 11, Page 1037) in the first line of Abstract and first line of Introduction, change "Beta vulgaris var. cicla)" to read "Brassica campestris L. Chinensis group)"
Corticium salmonicolor: 885
Corynebacterium michiganense: 336; 654
Corynespora cassiicola: 142
Cotton: estimated reduction in yield as result of parasitic diseases in 1970, 504; preharvest infection of cotton fiber by fungi in the U.S. Cotton Belt, 714
---: anthracnose, seedlings of several Gossypium species and varieties were screened for susceptibility to two isolates of Colletotrichum gossypii, 879
---: Aspergillus infection, living cotton fiber with typical bright greenish-yellow fluorescence (BGY) was regularly infected

with A. flavus, and incidence is centered in two areas of the U.S., 33
---: boll rot, from Aspergillus flavus, data indicate that high levels occur principally in certain geographically limited production areas in California, Arizona and Texas, 510
--- Fusarium wilt resistance, Auburn 56, Auburn M and Dixie King II varieties were highly resistant in a 4-year test, 896
---: nematodes, control of reniform nematode with systemic nematicides, 362
---: seedling disease: seed treatments with several nonmercurial fungicides appear equal to or better than the commonly used mercury alkyl treatments, 285; over-the-top sprays of fungicides for cotton seedlings, no treatment significantly better than lime water for control of disease, 307; several fungicides tested in vitro for their systemic activity and phytotoxicity on cotton seedlings, 1111
---: Verticillium wilt: bed and sidedress of benomyl-surfactant combinations gave greatest yield increase and delayed wilt symptom expression of Verticillium in field trials, 569; 90% of isolates of V. albo-atrum from wilt-infected olive trees in California were T-1 strain, the strain predominant in severely infected cotton from same area, 780; virulence of American and Australian isolates of Verticillium albo-atrum in cotton compared, 977
Cottonwood (see Populus deltoides)
Cover crop: a 1-year rotation of rye (nonhost of root-knot nematode) preceding a muskmelon crop in Indiana resulted in melon yields almost as high as after fumigation, 55
Cowpea: (Vigna sinensis)(southern pea): cowpea, pea, bean and Gomphrena globosa effective as selective hosts for identifying strains of alfalfa mosaic virus, 127; apparently first report of a new race of cowpea rust, Uromyces phaseoli var. vignae found in Georgia in August 1970, 384; anthracnose of cowpeas reached epiphytotic proportions at the University Farm, Ile-Ife, Nigeria, incitant tentatively identified as Colletotrichum lindemuthianum, 820
Cronartium comandrae: 697
--- fusiforme: 623
--- quercuum: 623
--- ribicola: 932
Crop residue decomposition: green crop residues amended to field or greenhouse soil for lettuce seedlings caused damage to their root systems, 837
Crown gall: of tomato, caused by Agrobacterium tumefaciens, retarded after bacteriophage treatment, 145
Crucifers: bok choy (Brassica campestris): bacterial soft rot and black spot diseases, 1037
---: broccoli and cauliflower (Brassica oleracea var. botrytis): four accessions of cauliflower of 148 tested showed resistance to Alternaria brassicicola, none of 9 broccoli lines showed resistance, 454; control of sugarbeet cyst nematode in broccoli with 1,3-D or DD nematicides, 1101; a firm head rot of broccoli found in California, Sclerotinia sclerotiorum and Erwinia carotovora consistently isolated, 1136
---: Brussels sprouts (Brassica oleracea var. gemmifera): basal sprout rot in 'Jade' is most directly associated with senescence; Alternaria tenuis and Fusarium roseum were fungi most often isolated from diseased plants, 297; imported varieties and cultivars of Brussels sprouts appear to be better hosts of sugarbeet nematode than local varieties in California; no differences observed for cabbage root nematode, 1005
---: cabbage (Brassica oleracea var. capitata), symptoms of Fusarium yellows were apparent simultaneously in both ozone and ozone-free filtered air chambers, 47; control of sting nematodes on direct-seeded cabbage, 412; preplant fumigation with nematicides significantly increased cabbage yields over check where Heterodera schachtii was present in high numbers, 841; controlled-atmosphere storage conditions and its influence on postharvest decay fungi of cabbage, 864; cabbage seed infection by Phoma lingam eradicated with 24-hr-soaks in suspensions of benomyl or thiabendazole, 934
---: downy mildew, measuring systemic infection in young hypocotyls of cauliflower, broccoli and cabbage, 877
---: turnip greens (Brassica rapa), fungicides significantly reduced leafspots and increased percentage of leafy greens acceptable for processing, 540
Cryptococcus fagi: 591
Cucumber (see Cucurbits)
Cucumber mosaic virus: on pepper, in North Carolina, 751
Cucurbits: 49 of 59 cucurbit species and cultivars tested were susceptible to bacterial wilt of cucurbits, the genus Cucurbita contained most of the resistant species, 816
---: cantaloup: benomyl better than hot water dip or approved chemicals, captan or chlorine, for control of stem scar and rind mold of cantaloup, 183

---: cucumber: angular leaf spot, caused by Pseudomonas lachrymans, was not easily seed-transmitted in California-grown seed, 3; greenhouse method developed for evaluation of cucumber varieties for resistance to Corynespora leaf spot disease, 142; pear virus, apparently first reported case of transmission of a virus from pear to an herbaceous host by pear psylla, 295; cucumber leaf was infected by the ash strain of tobacco ringspot virus transmitted by Xiphinema americanum, 475
---: muskmelons, response to fumigation for control of Meloidogyne incognita following 1 year of a nonhost crop of rye, 55
---: squash: presence of watermelon virus mosaic in New York and New Jersey, 132; cold pox of butternut squash was caused by low temperature and found in Collier County, Florida in November 1970, 350; all of 542 introductions of pumpkin and squash tested were susceptible to root-knot nematode, 666
Cupressus arizonica (Arizona cypress): causal fungus of the canker disease of Arizona cypress in South Carolina is Monochaetia unicornis, 810
--- lusitanica: root disease, caused by Poria vincta var. cinerea, found in East Africa for the first time, 257
Currant (Ribes nigrum): benomyl is effective for control of anthracnose, powdery mildew, and rust on black currants, 932
Curvularia eragrostidis: 38
--- cymbopogonis: 349
--- geniculata: 861
Cyclamen persicum: Septoria leaf spot was determined to be caused by Septoria cyclaminis, first report of disease in the United States, 93
Cylindrocladium sp.: 743
--- spp.: 419
--- crotalariae: 419
--- floridanum: 700; 1094
--- scoparium: 679
Cyprus: 707
Cytoplasm (see also Male-sterile cytoplasm): yield loss in corn with T cytoplasm compared with corn with N cytoplasm, 115; corn inbreds containing T cytoplasm were all susceptible to Helminthosporium maydis in Tennessee tests, 195; development of multiplasm so that cytoplasms may lessen likelihood of recurrence of disease problem brought about by single cytoplasm use in corn breeding, 400; seedling and mature plant resistance to Phyllosticta sp. in corn inbreds with normal and different male-sterile cytoplasms, 401; the importance of cytoplasmic inheritance in corn to susceptibility to H. maydis Race T and the possibility of similar effects in other crops with uniform cytoplasms such as grain sorghum, wheat, millet, sugarbeets, onions, pepper and petunia, 473; corn, seed tests of corn inbred F44 with normal and T cytoplasm indicate H. maydis and Fusarium moniliforme may be seed-transmitted especially in lines with T cytoplasm, 486; reactions of 5 corn inbred lines with different cytoplasms to H. maydis Race T toxin were determined in the field in New York, 938
Cytospora leucostoma: 314

Dactylium dendroides: 120
Dalbergia sissoo: hardwood tree severely attacked by Ganoderma lucidum in West Pakistan, 1130
Damping-off: of pea, Pythium aphanidermatum is cause of serious damping-off and root rot of pea in Iran, favored by high temperatures and excessive soil moisture, 244; of sugarbeets, caused by Aphanomyces cochlioides, was reduced in the greenhouse by several soil fungicides and soil amendments, 440
Defoliation: of bean by injury from rain, in New York, 457
Deposition: deposition and tenacity of foliage protectant fungicides on leaves of bush bean, soybean and cotton, 898
Desiccation: germinated conidia of Cercospora carthami and Ramularia carthami can withstand prolonged desiccation without loss of viability, 846
Diagnostic host plants: broad bean (Vicia faba), sweet basil, (Ocimum basilicum), and cockscomb (Celosia plumosa) found as new diagnostic hosts of papaya mosaic, 785
Diaporthe alleghaniensis: 788
--- phaseolorum var. sojae: 776
Didymella sp. (perfect stage of Phleospora idahoensis): found on overwintered culms of native fescues in Oregon, 63
Differentiation: of geranium plants from anther callus to produce virus-free plants, 1017
Diplodia natalensis: 149; 752
--- theobromae: 741
--- zeae: 158; 634
Dips: dipping raspberries in sodium salt of dehydroacetic acid (DHA-S) to extend shelf-life had adverse effect on flavor, 175; quick-dip method to inoculate tomato seedlings with tobacco mosaic virus, 375; warm fungicidal soaks for Fusarium basal rot of Iris improved control, 425; soft rot of sweetpotato controlled with dip treatments of DCNA, 1097

Dispersal: of spores of Scirrhia acicola, cause of brown-spot needle blight of long-leaf pine seedlings, 309
Distribution: of sugarbeet viruses transmitted by aphids in Washington, 103; of Claviceps gigantea on two lines of corn in Mexico shown in relation to temperature and precipitation, 231; of Dutch elm disease and the smaller European bark beetle in the United States as of 1970, 271; distribution and occurrence of Sclerotium oryzae on rice in California, 757; geographical distribution of Aspergillus flavus boll rot in experimental plots across the U.S. Cotton Belt in 1969, 33; in the U.S. cotton crop of 1970, 510; geographical distribution of certain preharvest fungal infection in cotton fiber in the U.S., in 1969-1970, 714; of Helminthosporium maydis Race T by States in 1969 and 1970, 302; distribution, race frequency, virulence, and mating type of isolates of H. maydis in the Northeastern United States in 1970, 495; of physiologic races of wheat stem rust in the U.S. in 1969, 187; of races of wheat stem rust in 1970 with relative prevalence, 986; of races of oat stem rust in the U.S. with relative prevalence, 992
Ditylenchus dipsaci: 433; 621
Dolichos biflorus: anthracnose, all accessions of D. biflorus seedlings tested showed extreme susceptibility to Colletotrichum dematium f.sp. truncata, 10
Dominican Republic: 207
Dothiorella ribis: 893
Dothistroma pini: 1040
Downy mildew: of sunflower, Plasmopara halstedii caused heavy damage in principal sunflower production areas of U.S., 11; corn inbreds tested for resistance to sorghum downy mildew, 202; of peas in northern Idaho and eastern Washington in 1970, 730; of lettuce, 51 plant introductions found immune to races 5 and 6 of downy mildew fungus, Bremia lactucae, 794; method for quantitative measurement of systemic infection of crucifers by Peronospora parasitica, 877
Dry bubble (Verticillium disease): of cultivated mushroom, control with benomyl, 684
Dutch elm disease (DED): treatments of American elm trees with benomyl by soil or trunk injections did not reduce symptoms of DED, 58; found on winged elm tree in Oconee County, South Carolina in September 1970, first report from this State, 158; soil injection of benomyl plus Surfactant F and benomyl plus Tween 20 reduced foliar symptoms of DED in elms treated prior to inoculation, 179; DED now found in all but 10 of the contiguous United States and the smaller European bark beetle has been reported from all but three, Florida, Arizona and Montana, as of 1970, 271; distribution in Minnesota in 1970, 682; relative resistance of diploid elm species to Ceratocystis ulmi, 980; spread of DED from New Plymouth to Mountain Home represents a long-range jump over arid treeless rangeland in Idaho, 1064
Dwarfing: exocortis virus associated with dwarfness in citrus trees in Australia, 998

East Africa: 257
Elderberry: several sap-transmissible viruses were isolated from foliage or roots of cultivated elderberry in New York, 913
Elm: addition of adjuvants to benomyl applied as a drench did not increase uptake of benomyl by American elm seedlings grown in planting media, 971
---: Dutch elm disease, 58; 158; 179; 271; 682; 980; 1064
---: elm phloem necrosis, found for first time in Pennsylvania in 1971, 1085
Endive (Cichorium endivia): plants with mottled leaves found to be infected by Bidens mottle virus, in Florida, 1061
Endothia sp.: 467
--- parasitica: 199; 201
England: 268
Entyloma sp: 207
Environmental modification: peppers resistant to potato virus Y in summer in Israel tests became susceptible under winter conditions, 604
Epicauta vittata: 628
Epidemic: sycamore anthracnose, in the Mississippi-Yazoo Delta in spring of 1970, caused by Gloeosporium nervisequum, 93; meteorological aspects of Cercospora epidemics on sugarbeets in the Texas-New Mexico High Plains, 1073
Epidemiology: of Leveillula taurica, in Iraq, 192
Epiphytotic: anthracnose of cowpeas in Nigeria, 820
Eradication: of chokecherry for control of X disease of peach, 645
Erwinia carotovora: 1037; 1136
--- tracheiphila: 816
Erysiphe graminis: 342
Etiological agents: of arecanut yellow leaf disease, 170
Eton-Bug disease: of sweet cherry, in Washington, thought to be of virus origin, 344
Eutypa armeniacae((imperf. Cytosporina): limb dieback of Ceanothus at pruning wounds was caused by pathogen responsi-

ble for apricot dieback, Eutypa armeniacae, 1006
Exposure chamber: with a contactless meter relay for controlling ozone concentrations in chamber, 774
Eyespot: of wheat, caused by Cercosporella herpotrichoides, controlled with benomyl, 45

Fall tillage: the host crops tobacco and corn had greater influence on nematode population density than fall tillage, 41
Fertilizer: a mineral fertilizer added in large amounts to nematode-infested soil reduced severity of tomato root knot, 1126
Field evaluation: for tomato early blight resistance, 807
Florida: 38; 139; 142; 154; 241; 261; 350; 390; 412; 447; 614; 848; 923; 929; 949; 1061
Flower blast: of statice (Limonium sinuatum) in south Florida, 848
Fluorescence: cotton fiber samples examined from 53 locations in U.S. to detect fluorescence associated with Aspergillus flavus boll rot, 33
Fomes annosus: 317; 490; 625; 659; 761; 1085
--- igniarius var. populinus: 662
Forking: of carrots, caused by Heterodera carotae, in Italy, 111; of carrots grown in muck soil, forking often accompanied "brown spot" damage, caused by Pythium spp., 536
France: 1090
Fungi: infection of corn seed with Helminthosporium maydis and other fungi in Indiana in 1970, 387; associated with southern pine needle blight, Lophodermella cerina, Ploioderma lethale and P. hedgcockii, 764
Fungicides (see also Antibiotics, Chemicals and Nematicides): nonmercurial fungicides for cottonseed were equal to or better than the alkyl mercury seed protectants, 285; home applications of garden dusts vary with the type of garden duster used, the pump gun type was superior to the plastic squeeze type, 1014
---: 2 AB (2-aminobutane), gave excellent control of Nectria decay of apple with postharvest chemical dip, 771
---: benomyl (methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate): controlled Verticillium wilt of tomato, 26; controlled eyespot lodging in wheat, 45; effective against black leaf streak of banana, 50; ineffective against Dutch elm disease, 58; prophylactic action against Dutch elm disease, 179; not effective in control of Heterodera avenae in wheat roots, 67; for control of brown rot of nectarines, 69; benomyl + ethephon effective against Fusarium corm rot of gladiolus, 82; on pearl millet, 87; showed selective fungitoxicity to fungi parasitic to commercial mushroom, 120; applied after washing to oranges and tangerines before degreening, 149; controlled strawberry anthracnose, 139; effective in control of a bark necrosis of Hevea brasiliensis, 152; effective against Fusarium blight on 'Merion' Kentucky bluegrass, 167; 333; inhibited growth of Endothia parasitica, in 2-year-old trees in the field, 199; best of fungicides tested for control of stem-scar and rind mold of cantaloup, 183; in heated suspensions for control of brown rot and Rhizopus rot of peaches, 228; effective against white and gray mold of beans, 234; controlled root rot of Poinsettia, 238; more effective than tribasic copper sulfate for control of sour orange scab on 'Page' citrus hybrid, 241; significantly less effective than Karathane for control of powdery mildew of apple, 263; equal to Karathane in control of apple powdery mildew, 514; as a spray or dip immunized chrysanthemum cuttings to infection by Verticillium albo-atrum and Rhizoctonia solani, 347; reduced disease severity of black root rot of tobacco, 352; as aerial low volume spray controlled Cercospora leaf spot of sugarbeet, 408; under sprinkler irrigation, 449; for control of Fusarium basal rot of Iris bulbs, 425; benomyl-surfactant combinations increased yield of cotton and delayed Verticillium wilt expression, 569; as a spray controlled Ramularia leaf spot on Primula polyantha, 586; gave good control of newly described blight of swordfern, 614; controlled Cylindrocladium scoparium of azaleas, 679; controlled Verticillium disease of cultivated mushroom, 684; effective against pecan scab at more extended interval applications than presently recommended materials; 711; applied as dust or dip controlled Fusarium tuber dry rot of potato, 723; preharvest sprays reduced decay of nectarines before and after harvest; postharvest treatment with DCNA (2,6-dichloro-4-nitroaniline) + benomyl in wax emulsion reduced decay development, 804; controlled apple scab, 828; incorporated into green crop amended soil reduced root damage of lettuce seedlings, 837; time of application in relation to control of Botrytis strawberry fruit rot, 883; as soil application gave best control for Verticillium wilt of tobacco, 889; effective for control of leaf diseases on black currant, 932; completely eradicated Phoma lingam in infected cabbage seed with a 24-hr seed soak, 934; uptake of

benomyl by American elm seedlings grown in planting media was not improved with adjuvants, 971; late fall treatment of strawberry plants with benomyl controlled storage decay, 1009; effective against stem canker of tea plant, 1021; as a slurry seed dressing controlled loose smut of wheat (Ustilago tritici), 1053; for cotton seedling disease control, 1111; controlled Cercospora leaf spot of Phlox, 1120, effective as seed treatments for common bunt of wheat, 1132
---: Bordeaux, and cuprous oxide superior to sulfur fungicides and organics against coffee rust in the Philippines, 216
---: Busan 72 (see TCMTB)
---: captafol (see Difolatan)
---: captan: for seed treatment of pea against Pythium, 244; for cottonseed treatment, 285; for black root rot of tobacco, 352; seed treatment for barley and oats, 1122
---: carboxin (Vitavax): seed treatments to control: pearl millet against smut, 87; cotton diseases, 285; head smut of corn, 501; dwarf bunt of wheat, 613; Sclerotium root rot of sugarbeet, 630; loose smut of wheat, 1053; cotton seedling diseases, 1111; barley and oat diseases, 1122; common bunt of wheat, 1132
---: Cela W 524, a new systemic fungicide was highly effective against wheat leaf rust in a growth chamber, 691
---: Ceresan M-DB, treatment of barley seeds did not reduce viability during first 20 months of storage, anaerobic and hot water treatments did, 118
---: chloroneb: reduced sugarbeet root rot, 630; for cotton seedling disease control, 1111; controlled Typhula blight of creeping bentgrass caused by Typhula itoana, 1118; controlled barley leaf stripe, 1122; chloroneb and chlorothalonil in over-the-top sprays for cotton seedlings gave significantly better stands than the "wet" control, but not significantly better than lime water, experiment also pointed up the fallacy of comparing treatments with aqueous sprays with a "dry" control, 307
---: chlorothalonil: 307; controlled Rhizoctonia blight of swordfern, 614
---: copper fungicides: for control of coffee rust in Kenya and Brazil, 209; cuprous oxide for coffee rust in the Philippines, 216; copper fungicides reduced leafkill by Cercospora beticola on sugarbeet, 479; cupric hydroxide in greenhouse trials as foliage spray controlled Pseudomonas syringae on beans, 735; neutral copper for pink pitting of grapefruit, 923
---: DCNA: controlled Rhizopus rot of peaches, 228; controlled white and gray mold of beans, 234; preharvest benomyl spray and postharvest wax emulsion with DCNA controlled decay of nectarines, 804; treatment of root pieces of sweetpotatoes prior to prespouting reduced incidence of soft rot from Rhizopus sp., 1097
---: Dexon: Dexon, Difolatan and maneb most promising of chemicals tested in soil dilution plates for injection into soil to prevent crown rot of apple, 213; effective in reduction of damping-off of sugarbeets in greenhouse soil, 440; reduced root damage to lettuce seedlings in crop amended soil, 837
---: DHA-S (sodium salt of dehydroacetic acid), sprayed on raspberries before harvest extended the shelf-life of fruits, 175
---: 1,3-D (1,3-dichloropropene): 1,3-D + chloropicrin, more effective than 1,3-D alone in suppressing Verticillium albo-atrum infection in potato stems in Connecticut, 968
---: Difolatan: effective against bark necrosis of Hevea, 152; tested in soil dilution plates for injection into soil to prevent crown rot of apple, 213
---: dithiocarbamate, in an oil-in-water emulsion applied to bananas by aircraft controls Sigatoka and black leaf streak in Central America and the Philippines, 437
---: DMTT, controlled black root rot of tobacco and bean, 352
---: Dyrene, and zinc ion + maneb controlled Helminthosporium sorokinianum on Kentucky bluegrass, 959
---: ETMT: proved superior to Dexon in control of Phytophthora and Pythium in bedding plants, 587; controlled Pythium root rot disease of potted chrysanthemums, 851; for cotton seedlings, 1111; ETMT + PCNB, 1122
---: ferbam, dip at 55°C for 1/2 min gave complete control of sweetpotato scurf, 746
---: Geigy 20072, for control of decay of degreened tangerines, 149
---: Karathane: significantly better than benomyl against powdery mildew of apple in Oregon when evaluated over an entire season, 263; benomyl and triarimol equal to Karathane for control of powdery mildew of apple in Washington, 514
---: maneb: for possible soil injection against crown rot of apple, 213; applied weekly gave control of Stemphylium leaf spot of garden and sultan balsam, 374; effective in control of leaf spots of turnip greens for processing market, 540; controlled Botrytis leaf blight of onion in New York,

565; maneb and chlorothalonil sprays no more effective in controlling Zoysia rust and St. Augustine blast on lawn grass than prompt removal of mower clippings, 778; against seedborne rust of safflower, 1046; for control of needlecast fungus, Lophodermium pinastri, 1116
---: MENCS, controlled black root rot of tobacco and bean, 352
---: mercurous chloride, (60%) and 30% mercuric chloride granular formulation effective against Typhula blight when applied 1 month before snowfall, 1118
---: methyl bromide: best of three fumigants tested for control of Fusarium wilt of petunia, 580; preplant fumigation with methyl bromide + chloropicrin to kill Phytophthora parasitica in citrus groves, 929
---: NF-35 (thiophanate): controlled Fusarium blight of 'Merion' Kentucky bluegrass, 333; controlled apple scab, 828
---: NF-44 (thiophante-methyl), controlled apple scab, 828
---: organo-mercury, controlled bark necrosis of Hevea, 152
---: oxycarboxin (Plantvax), was better than carboxin, benomyl, ziram for control of smut of pearl millet in India, 87
---: PCNB, and benomyl controlled stem and stolon infection of seed potatoes by Rhizoctonia solani when used as soil treatments, 550
---: Plantvax (see oxycarboxin)
---: PMAA, for seed treatment of barley and oats, 1122
---: potassium azide: fungicidal activity as affected by pH, 1077; controlled microsclerotia of Cylindrocladium floridanum in artificially infested soil, 1094; effective seed treatment, 1122
---: SMDC, controlled black root rot of tobacco and bean, 352
---: sodium azide, effective seed treatment for barley and oats, 1122
---: SOPP, for decay of degreened oranges and tangerines, 149
---: TCMTB (Busan 72), as cottonseed treatment, 285; effective against seedborne rust of safflower, 1046
---: Terracoat L21, cottonseed treatment, 285
---: tetrachloroisophthalonitrile, controlled tan leaf spot of Rhoeo discolor, 38
---: thiabendazole: materials for a field test kit to test concentration of thiabendazole in treating tanks for control of bulb diseases, 77; gave control for Fusarium rot of gladiolus corms, 82; mushroom diseases, 120; decay of degreened oranges and tangerines, 149; white and gray mold of beans, 234; black root rot of tobacco and bean, 352; Fusarium basal rot of iris bulbs, 425; both thiabendazole and benomyl are superior substitutes for mercurials in control of Fusarium basal rot of Narcissus, 428; controlled leaf spots of turnip greens, 540; control of dwarf bunt of wheat, 613; blight of swordfern, 614; thiabendazole, not effective as bed drench or cutting soak for control of Cylindrocladium rot of azalea, 679; controlled Fusarium tuber dry rot of potato, 723; effective as preinoculation treatment for control of Diplodia rot of mango fruits, 752; extremely effective in control of decay of mechanically harvested 'McIntosh' apples, 791; eradicated Phoma lingam from cabbage seed, 934; controlled Cercospora leaf spot of Phlox, 1120
---: thiophanate (see NF-35)
---: thiophanate-methyl (see NF-44)
---: thiram: and captan seed treatment of pea in Iran, preliminary results indicate seed treatments control Pythium aphanidermatum and thus increase yields, 244; cottonseed treatment, 285; for oat and barley seeds, 1122
---: triarimol equal to Karathane for control of apple powdery mildew, 514
---: Vitavax (see carboxin)
---: zinc ion and maneb; showed promise for control of tan leaf spot of Rhoeo discolor, 38; controlled Fusarium tuber rot of potato, 723; controlled Helminthosporium sorokinianum on Kentucky bluegrass, 959; for seed treatments of barley and oats, 1122
---: zineb: for control of Septoria leaf spot of pistachio nut, 72; 1.3% oil, zineb, or neutral copper sprayed first part of July controlled pink pitting of grapefruit in Florida, 923
---: ziram, not effective against smut of pearl millet, 87
---: several tested together: for control of bean root rots, 483; for control of the sweetpotato black rot fungus, Ceratocystis fimbriata, 523; 19 commercial fungicides tested for their deposition and persistence at fungistatic concentrations on the leaves of bush bean, soybean and cotton plants, 898; for effectiveness in control of cotton seedling diseases, 1111
Fusarium sp.: 580; Fusarium wilt of tomato, announcement of monograph, Fusarium wilt of tomato, 653; 788; 1124
--- spp.: 337; 714
--- moniliforme: 486; 634; 687; 1046
--- oxysporum: 687
--- --- f. sp. cepae: 1000
--- --- f. sp. conglutinans: 47

--- --- f. sp. gladioli: 82; 340; 425
--- --- f. sp. narcissi: 428
--- --- f. sp. pisi: 1032
--- --- f. vasinfectum: 896
--- roseum: 167; 297; 333; 687; 861; 864;
--- --- sambucinum: 723
--- solani: 152; 390; 687; 861
--- --- f. phaseoli: 483
Fusicladium dendriticum: 509
--- effusum: 711

Ganoderma lucidum: 1130
Garden balsam (Impatiens balsamina): Stemphylium leaf spot, found in greenhouse in New Jersey, 374
Garden dusters: pump gun garden dusters proved more uniform in amounts of dust expelled than plastic squeeze dusters and did not exceed approved dust levels, 1014
Gardenia: nematicide-insecticide D-1410 was as effective as cynem and BAY 68138 for control of Meloidogyne incognita, 357
Genetic stability: of urediospores of wheat stem rust in cryogenic storage, 695
Genotypes: of corn and their possible role of improving insight into the nature of the nuclear and cytoplasmic interactions which control resistance to Helminthosporium maydis Race T, 938
Georgia: 10; 312; 384; 453; 540; 623; 741; 951
Geranium (Pelargonium hortorum): differentiation of virus-symptomless geranium plants from anther callus, 1017
Germ plasm: of red clover, with high resistance to northern anthracnose, 920
Gibberella zeae: 634
Gladiolus: corm rot, corm soaks of heated water with either benomyl or thiabendazole + ethephon resulted in production of larger and heavier corms, 82; CO_2 treatment of gladiolus corms was very effective in revealing latent Fusarium infections, 340
Gloeosporium sp.: 424
--- nervisequum: 93
Glume blotch: of wheat, in India, Phoma insidiosa found to cause glume blotch for first time, 95
Gomphrena globosa: one of four selective hosts utilized to identify strains of alfalfa mosaic virus, 127
Gossypium (see Cotton)
Grafting: of tomato on to resistant rootstocks to control tomato bacterial wilt, 521
Graminella nigrifrons: 273
Grape: Pierce's disease, suppressed in vines and rooted cuttings with soil drenching of tetracycline, 610; a bark and wood pitting disease found in California, 860; a stem pitting disease of grapevine was observed in Greece for the first time in 1970, 869
Grasses: 52 isolates of Cochliobolus carbonum from 20 grass genera were tested for pathogenicity to 22 gramineous species, and showed lack of host specificity, 325
---: beachgrass, American, new disease caused by Marasmius species found in this important beach grass, in North Carolina, 582
---: creeping bentgrass, comparison of application dates for control of Typhula blight, 1118
---: fescue (Festuca spp.). Phleospora idahoensis, cause of stem eyespot, found for the first time in Oregon in 1970, 63
---: johnsongrass, host of a strain of corn stunt disease in South Carolina, 273
---: 'Merion' Kentucky bluegrass: Fusarium blight controlled with benomyl drench, 167; systemic fungicides benomyl and thiophanate effectively controlled Fusarium blight, 333; excised blades used for fungicidal evaluation against Helminthosporium sorokinianum, 959
---: St. Augustinegrass, and zoysiagrass, removal of mower clippings from lawn in Mississippi effectively controlled Puccinia zoysiae and Pyricularia grisea, without fungicides, 778
Gray mold: of bean, caused by Botrytis cinerea, field control on beans in Washington with DCNA, and three benzimidazoles, 234
Greece: 869; 1049
Greening disorder: of Trillium grandiflorum, 1108
Guatemala: 207
Guignardia aesculi (Phyllosticta sphaeropsoidea): 37

Halo blight (see Blight)
Hardwoods: Ganoderma lucidum, a virulent incitant of basal stem rot of hardwoods in West Pakistan, 1130
Hawaii: 7; 392; 521; 743; 783; 785; 941
Head mold: of grain sorghum, in Kentucky, was caused principally by Fusarium spp., 337
Helianthus annuus (sunflower): outbreak of downy mildew, 11; a vascular wilt-inciting Fusarium isolated from diseased sunflower in Texas, first report, 1124
Helicotylenchus multicinctus: 707
Helminthosporium sp. (Cochliobolus carbonum): 325
--- gramineum: 858; 1122
--- maydis: 99; 115; 195; 302; 371; 382; 387; 392; 469; 473; 486; 495; 563; 667; 672; 726; 755; 759; 855; 938; 947; 949; 951; 954; 956; 966
--- sigmoideum: 498

--- sorokinianum: 959
Hemileia vastatrix: 209; 216
Hemlock (Conium maculatum): western celery mosaic virus, hemlock was the most frequent host of the virus found near celery crops in England, 268
Hendersonia acicola: 764
Heracleum sphondylium (hogweed): weed host of western celery mosaic virus in Britain, 268
Herbicides (see Chemicals)
Heterodera spp.: 399
--- avenae: 67
--- carotae: 111
--- cruciferae: 1005
--- glycines: 13
--- lespedezae: 114; 927; 974
--- schachtii: 841; 1005; 1101
Hevea brasiliensis (para rubber): new disease of para rubber, caused by Fusarium solani and Botryodiplodia theobromae, 152
High-lysine corn: extreme susceptibility to kernel and ear rots evident in 1971 when demand for "normal" cytoplasm corn resulted in increased planting of high-lysine corn, 1046
Histology: of Phoma lingam-infected cabbage seed, and control of the pathogen, 934
Holly (Ilex cornuta var. burfordi): Gloeosporium canker found on Burford holly in Alabama nursery, apparently for the first time, 424
Honduras: 207; 415
Hoplocampa testudinea: 828
Horse hair blight: of cacao, caused by two Marasmius species, in the Solomon Islands, 885
Host range: of Radopholus similis from banana areas of Central America, 415; of Erwinia tracheiphila on members of the Cucurbitaceae, 816
Hot water treatments (see Temperature)
Hypoxylon mammatum: 650; 662

Idaho: 54; 501; 550; 730; 1064
Illinois: 37; 96; 114; 573
Immunity: to downy mildew (Bremia lactucae) in lettuce introductions, 794
India: 87; 95; 107; 170; 185; 283; 365; 526; 544; 593; 630
Indiana: 55; 387
Indicator: evaluation of Shirofugen as a reliable indicator for rose mosaic virus, 431
Inheritance: tomato anthracnose resistance in PI 272636, 253
Inoculation: tomato bacterial wilt, was produced in susceptible scions on resistant rootstocks only by introduction of the bacterium through stem wounds, 521; artificial inoculation of strawberry varieties to test for resistance to Mycosphaerella fragariae, 573; the leaf water-soaking and stem inoculation for comparison of disease reaction of bean varieties to Xanthomonas phaseoli and X. phaseoli var. fuscans bacteria, 577; inoculation of mushrooms with Verticillium malthousei spores earlier than 21 days after casing caused lowest mushroom yield and highest disease incidence, 643; cross-inoculation studies give proof of Cercospora microsora and C. exitiosa identity, 733
Insecticides (see Chemicals)
In vitro: selective fungitoxicity of benomyl and thiabendazole against parasitic fungi of commercial mushroom could be demonstrated, 120; evaluation of fungicides for control of Helminthosporium sorokinianum with use of excised blades of Kentucky bluegrass, 959; benomyl showed toxicity to Phomopsis theae, cause of tea stem canker, 1021; use of cotton seedlings in vitro for measuring comparative systemic fungicidal activity and phytotoxicity of certain seed and soil fungicides for control of cotton seedling diseases, 1111
Iowa: 317; 479; 855
Iran: 85; 244
Iraq: 192
Ireland: 175
Iris: Fusarium basal rot of bulbs, immediate treatment after harvest, 30 minute soak in benomyl or thiabendazole at 40°C and addition of Biofilm improve control, 425
Isolates: pathogenic variation in Philippine isolates of Xanthomonas oryzae to 24 rice varieties, 22; pathogenic variability in Indian isolates of X. oryzae, 593; of Helminthosporium maydis, overwintering and survival capacities, 99; the virulence and mating types of H. maydis collected in 1970, 759; Phytophthora parasitica isolate from white clover inoculated on four soybean varieties to test pathogenicity, 130; Cochliobolus carbonum isolates from 20 grass genera showed marked variation in pathogenicity but a lack of host specificity, 325; of American and Australian strains of Verticillium albo-atrum, 977; of mechanically transmissible virus from sorghum, in France, 1090
Israel: 408; 604
Italy: 111
Ixora coccinea: tropical shrub in Florida susceptible to bacterial leaf spot, caused by Xanthomonas maculifoliigardeniae, 261

Juniperus virginiana (Eastern redcedar): plants wound-inoculated with Monochaetia unicornis developed cankers but no can-

kers were found on redcedar in South Carolina survey, 810

Kabatiella caulivora: 920
Katsura-tree (Cercidiphyllum japonicum): Dothiorella ribis associated with cankers on this ornamental tree at Brooklyn Botanic Garden, 893
Kentucky: 337
Kenya: 802
Kernel infection: of corn produced in 1970, with Helminthosporium maydis, 726

Latent infections: of Fusarium oxysporum f. sp. gladioli, rotting triggered with CO_2 corm treatment, 340
Leaf blotch: of Aesculus, additional species susceptible to blotch infection, caused by Guignardia aesculi, 37
Leafhopper: control of leafhopper-borne plant diseases possibly due to direct action of systemic biocides on mycoplasmas, 291
Leaf scorch: of London plane trees, no apparent correlation between nematode infestation and scorched symptoms on leaves, 280; of peanut, caused by Leptosphaerulina crassiasca, genetic constitution of plant seemed to condition the susceptibility to the pathogen and its expression as leaf scorch or pepper spot, 530
Leaf spot: of banana, 437; of broccoli and cauliflower, some resistance found in four cauliflower introductions, 454; of cucumber, angular leaf spot, caused by Pseudomonas lachrymans, 3; cucumber varieties tested for resistance to Corynespora cassiicola, 142; of garden and sultan balsam, caused by Stemphylium botryosum, 374; of Ixora coccinea, caused by Xanthomonas maculifoliigardeniae, 261; Manilkara hexandra, caused by Pestalotia palmarum, 185; of pistachio, caused by Septoria pistaciarum, in Texas, 72; of hybrid primrose, caused by Ramularia primulae, 586; of safflower, caused by Cercospora carthami and Ramularia carthami, 846; of sugarbeet, control of Cercospora beticola, 408; control of C. beticola on sugarbeet with fungicides, 479; Cercospora leaf spot epidemics on sugarbeets, 1073; of turnip greens, leaf spot control with fungicides significantly increased yield of leaves for processing, 540
Leaf stripe: bacterial, of Strelitzia reginae, 447
Legumes: cyst nematode, 18 species of legumes tested for suitability as hosts for Heterodera lespedezae, 927
Lepidium virginicum (peppergrass): weed host of Sclerotinia sp. in alfalfa fields in Maryland, 1087
Leptosphaerulina crassiasca: 530
Lespedeza: lespedeza cyst nematode, Heterodera lespedezae found in Illinois on lespedeza cultivar Kobe, 114; host of H. lespedezae, 927
Lethal decline disease: of strawberry, in Oregon, first reported occurrence, in State, 560
Lettuce: downy mildew, sources of immunity to Bremia lactucae found in over 50 plant introductions, 794; damage of lettuce seedlings related to crop residue decomposition, 837; natural infection of lettuce and endive by Bidens mottle virus in Florida, 1061
Leveillula taurica: first known report of occurrence in Iraq on sunflower and tobacco, 192
Limb dieback: of Ceanothus, by Eutypa armeniacae, apricot dieback pathogen, 1006
Lime-tree (Tilia platyphyllos): cross-inoculations proved that leaf-spotting (Cercospora microsora) fungus can attack stems and shoots and stem parasite (C. exitiosa) can attack leaves of lime-tree, 733
Linked genes: for resistance to soybean cyst nematode and reniform nematode in soybean is suggested, 1043
Lodging: lodging of wheat due to eyespot, controlled with benomyl, 45
London plane tree (Platanus x acerifolia): new, undescribed species of Meloidogyne sp. caused galled roots on London plane trees in Washington, D.C. area, 280
Longevity (see Survival and Viability)
Longidorus elongatus: first report of occurrence in British Columbia, 266
Lophodermella cerina: first report of L. cerina on loblolly and slash pines, 764
Lophodermium sp.: 309
--- australe: 764
--- pinastri: 1116

Macrosiphum avenae: 874
Malaysia: 152
Male-sterile cytoplasm (see also Cytoplasm): infection of corn kernels with Tms cytoplasm by Helminthosporium maydis in Minnesota seed tests, 371
Mango (Mangifera indica): preinoculation treatments with thiabendazole completely controlled Diplodia rot of mango fruits and oils controlled Rhizopus rot up to 20 days, 752
Manilkara hexandra: found to be host of Pestalotia palmarum, in south India, 185

Manitoba: 797
Marah oreganus (wild cucumber): wild cucumber mosaic virus, first report of this virus in M. oreganus in Oregon, 456
Marasmius sp.: 582
--- equierinis: 885
--- pulcher: 885
--- sarmentosus: 885
--- scandens: 885
Market diseases: of bok choy (Brassica campestris), 1037
Marmor lacerans: 645
Maryland: 280; 379; 755
Massachusetts: 687
Mating types: of Helminthosporium maydis isolates collected in 1970 in the Northeastern United States, 495
Meloidodera charis: 115
Meloidogyne sp.: 61; 280
--- hapla: 89; 1069
--- incognita: 41; 55; 357; 666; 1082; 1126
--- javanica: 607; 707; 1082
Methods and techniques: a procedure for estimation of concentrations of thiabendazole from 600 to 2000 ppm in treating tanks for control of bulb diseases, materials may be assembled into a field test kit, 77; artificial system for differentiating strains of alfalfa mosaic virus, 127; multiplasm, a proposed method for the utilization of cytoplasm in pest control, 400; modified water culture technique for use in growing and screening strawberry seedlings for resistance to Phytophthora fragariae, 471; use of competitive organisms in chain oil of power saw to reduce buildup of forest root rot pathogens in stumps, 659; a quantitative procedure developed for measuring mycelial growth of Peronospora parasitica in young hypocotyls of cauliflower, broccoli and cabbage, 877; freezing of conidia of Venturia inaequalis for storing inoculum 976
---: equipment: sediment trap used to collect dust samples to determine extent of nematode movement by wind, 61; sampling methods for oak wilt compared, the hammer and branch methods appear to be equally effective, 281; device for controlling ozone concentration in an exposure chamber, 774; a rotating-drum spore trap using Scotch tape as a trapping, staining and mounting substrate, for airborne spores, 832
---: inoculation: artificial inoculation of rice varieties for resistance to Xanthomonas oryzae isolates, 17; a technique of inoculating X. translucens f. sp. oryzicola on rice with sandpaper and sand rubber plates gave better openings for bacteria than punctures made with needles, 404; inoculation of tomato seedlings with tobacco mosaic virus by a dipping method was compared with a rubbing method and found effective and less time consuming, 375; rapid screening techniques (by injection or spray) of toxin solutions of monoconidial isolates of Helminthosporium maydis, Race T for selection of resistant corn plants, 469; excised-petiole inoculation for recovery of Corynebacterium michiganense from infected overwintered stems of tomato, 654; use of seedling injection to identify four bean blight bacteria, 703
Mexico: 231
Michigan: 399; 651; 788; 961; 1108; 1118;
Microsphaera grossulariae: 932
Minnesota: 11; 371; 498; 650; 682; 1026; 1040
Mississippi: 93; 309; 460; 778; 954
Missouri: 1085
Moisture: excessive soil moisture and high temperature favor development of damping-off and root rot of pea in Iran, 244; relation of high moisture, temperature, and RH in the development of water congestion of pea in Wisconsin, 249
Monilinia fructicola: 69; 228; 804; 898
--- laxa: 69
--- vaccinii-corymbosi: 651
Monilochaetes infuscans: 746
Monochaetia mali: 96
--- unicornis: 810
Morus alba: hardwood tree of West Pakistan infected by Ganoderma lucidum developed basal stem rot, 1130
Mower clippings: prompt removal from St. Augustine and zoysiagrass lawns controlled St. Augustine blast or gray spot and Zoysia rust in Mississippi, 778
Multiplasm: a multiplasm approach to plant breeding is proposed as a safeguard against a monocytoplasm approach (as a male sterile) to prevent recurrence of an outbreak such as southern corn leaf blight, 400
Mummy berry disease: of blueberry, 13 varieties and 47 selections tested for resistance; E-176 only selection resistant to both ascospore and conidial infection, 651
Mushroom (Agaricus bisporus): diseases, control of Mycogone perniciosa, Verticillium malthousei, and Dactylium dendroides with benomyl and thiabendazole, 120; effect of time of inoculation on incidence and control of Verticillium disease of commercial mushroom, 643; control of Verticillium disease of cultivated mushrooms with benomyl spray applications to cased trays, 684
Mycogone perniciosa: 120

Mycoplasma: possibility of mycoplasma as etiological agents of arecanut yellow leaf disease, in India, 170; hypothesis proposed that systemic organo-phosphorus and oxime carbamate insecticides act directly on the causal organism of tobacco yellow dwarf aside from vector control, 291; suppression of Pierce's disease of grape with tetracycline suggests disease may be caused by a mycoplasma-like organism, 610; mycoplasma like bodies found in sprouting onion bulbs in storage, 824; mycoplasma-like bodies found associated with a flower greening disorder of a wild flower, Trillium grandiflorum, 1108
Mycorrhizal fungus: Rhizoctonia fragariae, fungus is a primary incitant of black root rot of strawberry under certain environmental and nutritional conditions, 599
Mycosphaerella sp.: 50
--- fragariae (Ramularia tulasnei): 573
--- musicola: 437
Myzus persicae: 526; 676

Narcissus: soak of 30 minutes at 25°C in benomyl or thiabendazole immediately after digging Narcissus bulbs is superior replacement for mercurial PMA in control of Fusarium basal rot, 428
Nebraska: 115; 479
Nectarine: brown rot, postharvest treatment with benomyl generally better than benomyl preharvest sprays in reduction of brown rot, 69; preharvest benomyl sprays and postharvest wax-fungicide treatments for control of decay of California nectarines, 804
Nectria coccinea var. faginata: 591
--- galligena: 445; 662; 771
Needle blight (see Blight)
Needlecast: in Scotch pine Christmas tree plantations, a new disease problem, 1116
Nematicides (see also Antibiotics, Chemicals and Fungicides): aldicarb, controlled lesion nematodes on snapdragons and chrysanthemums, 276
---: BAY 68138: effective as a bare-root dip on tomato and cabbage transplants against soybean cyst nematode, 13; bare-root dips protected citrus seedlings from Radopholus similis and Tylenchulus semipenetrans, 154; controlled lesion nematodes on snapdragons and chrysanthemums in heavy greenhouse soils, DBCP and chloropicrin did not, 276
---: BAY 25141: controlled soybean cyst nematode used as bare-root-dip on tomato and cabbage, 13; controlled nematodes on citrus seedlings, 154; compared with D-1410, 357
---: cynem: controlled Heterodera glycines on transplants, 13; control of nematodes on citrus seedlings, 154; compared with D-1410, 357
---: DBCP: after a muskmelon crop in Indiana, a season of rye (nonhost of Meloidogyne incognita) apparently reduced the need for fumigation before muskmelon planting, 55; applied in May and November ber in Cyprus for control of nematodes in banana plantations, 707
---: 1,3-D: 1,3-D, methyl bromide + chloropicrin, and gel as preplant fumigants increased cabbage yields, 841; 1,3-D or 1,3-D + chloropicrin against Verticillium albo-atrum, 968; 1,3-D and DD controlled Heterodera schachtii, 1101
---: DD: used to test effect of Pratylenchus penetrans on Verticillium wilt of tomato, 673; controlled Heterodera schachtii, 1101
---: Dupont 1410 (D-1410): nematicide-insecticide, gave best control of several nematicides on field-grown roses, to control Meloidogyne hapla, 89; compares favorably with cynem and BAY 68138 for control of M. incognita on gardenia, 357; controlled Pratylenchus penetrans in roots of apple, peach, pear, and cherry by foliar sprays, 617; D-1410 and DD applied to soil to test the effect of P. penetrans on Verticillium wilt of tomato, 673
---: several tested together: controlled Verticillium wilt of tomato up to 3 months, 26; to control Meloidogyne hapla, 89; several new systemic nematicides effective in control of Rotylenchulus reniformis in cotton, 362; controlled sting nematodes in direct-seeded cabbage in Florida, 412
Nematodes: estimate of cotton loss from nematodes by State, 504
---: burrowing: and citrus nematodes, Radopholus similis and Tylenchulus semipenetrans, controlled on citrus seedlings by chemical dips, 154; host range of Radopholus similis from banana areas of Central America with indications of additional races, 415
--- cyst: cereal cyst nematode, invasion of wheat roots by Heterodera avenae was not suppressed by benomyl when mixed with soil or as a root soak, 67
---: ---: grass, Heterodera punctata, first confirmed report of occurrence in Michigan; H. schactii, H. weissi, and H. trifolii also present, 399
---: ---: Heterodera carotae causes alterations in growth and shape of carrots infested with this cyst nematode in Italy, 111
---: ---: lespedeza, did not reproduce in

eight soybean differentials, 974; in Illinois, 114
---: ---: soybean, nematicides BAY 68138, BAY 25141 used as bare-root dips gave control with no phytotoxic effects, 13; the possible relationship in soybean of gene linkage for resistance to soybean cyst nematode and reniform nematode is thought to occur, 1043
---: ---: sugarbeet, preplant fumigation for Heterodera schachtii increased yields of cabbage significantly, 841; cabbage root nematodes, their reproduction on cultivars of Brussels sprouts, 1005; depth of application of 1,3-D or DD and efficacy in control of Heterodera schachtii on sugarbeet and broccoli in California, 1101
---: cystoid: Meloidodera charis recovered from corn field in Nebraska, 115
---: dagger: Xiphinema americanum can apparently act as a soilborne vector of the ash strain of tobacco ringspot virus, 475
---: lesion: Pratylenchus penetrans, controlled in heavy soils in greenhouse with aldicarb and BAY 68138 for at least 10 months, 276; effect of P. penetrans on the resistance of a pea variety to Fusarium wilt, 1032
---: needle: Longidorus elongatus damaging to strawberry plants in British Columbia, 266
---: plant parasitic, several nematode genera found associated with London plane tree but no correlation is apparent between their occurrence and scorch symptoms on leaves, 280
---: reniform: Rotylenchulus reniformis on cotton controlled with nematicides, 362; resistance to R. reniformis found in several soybean breeding lines, gene linkage for resistance to soybean cyst and reniform nematodes is suggested, 1043
---: root-knot: larvae of Meloidogyne sp. were among 28 genera recovered from dust traps in west Texas, wind could be important means of dissemination, 61; new species of Meloidogyne associated with galled roots of London plane tree in Washington, D.C., 280; M. hapla eradicated from field-grown roses with D-1410, 89; sainfoin was less tolerant than alfalfa to M. hapla at all ages in tests in Utah, 1069; M. incognita and Pratylenchus zeae population increase is influenced more by crop host than by fall tillage, 41; planting of a nonhost crop (rye) before planting melons appears to lessen need for fumigation against M. incognita infestation, 55; M. incognita controlled on gardenia with nematicides D-1410, cynem and BAY 68138, 357; 542 introductions of Cucurbita spp. including C. pepo, C. maxima and C. moschata all proved susceptible to M. incognita, 666; M. incognita suppressed in soil amended with oat straw and mineral fertilizer, 1126; 35 crops tested for resistance to root-knot nematodes, M. incognita and M. javanica, 1082; occurrence and reproduction of M. javanica on three citrus species in California, 607; DBCP fumigation after planting bananas reduced M. javanica and spiral nematodes in roots and surrounding soil in Cyprus, 707
---: root lesion, no correlation between Pratylenchus penetrans and incidence of Verticillium wilt of tomato was found in Ohio tests, 673
---: spiral, Helicotylenchus multicinctus, 707
---: stem: Ditylenchus dipsaci, causal agent of "tulip root" of oats in the coastal areas of California; 'Curt' was found to be tolerant in test trials, 'Sierra' and 'Kanota' were very susceptible, 433; nematode-inoculated alfalfa seedlings showed reduced numbers when grown for 24 days in presence of dichlorvos pest strips, 621
---: sting, several nematicides effective in control of Belonolaimus longicaudatus on direct-seeded cabbage in Florida, 412
New diseases: anthracnose on Dolichos biflorus in Georgia, 10; a malady of 'Jubilee' sweet corn reported from western Washington, 57; a bark necrosis of Hevea brasiliensis found to be caused by Fusarium solani and Botryodiplodia theobromae, in Malaysia, 152; leaf disease of bean, caused by Entyloma sp., found in Guatemala, Honduras and the Dominican Republic, 207; Marasmius blight of American beachgrass in North Carolina, 582; new petal blight on camellia caused by Pestalotia sp., 697; collar rot of papaya caused by Calonectria sp., 743; oleander blight caused by Pseudomonas syringae, in California, 1030; of broccoli, head rot, 1136
New distribution: a virus similar to sorghum red stripe and the johnsongrass strain of maize dwarf mosaic virus has been found in France, 1090
---: alfalfa mosaic virus was found in two previously unreported hosts, parsley and carrot, in California, 322
---: angular leaf spot of strawberry in Florida, 142
---: apple twig canker, caused by Monochaetia mali found in Illinois, 96
---: bacterial canker of sweet cherry identified

in Michigan, 961
---: the black rot organism, Ceratocystis adiposa, was isolated for first time from sugarcane cuttings in Hawaii, 7
---: Colletotrichum graminicola cause of heavy infection on corn, apparently for first time in Maryland, 755
---: Curvularia cymbopogonis found on rice seed samples from the Philippines, 349
---: Dothistroma pini cause of damage to Austrian pine in Minnesota, 1040
---: of Dutch elm disease, first report of the disease in South Carolina, 158
---: first record of elm phloem necrosis in Pennsylvania, 1085
---: Fomes annosus found on jack pine in Lucas County, Iowa, believed to be the first occurrence, 317
---: head smut of sorghum, caused by Sphacelotheca reiliana, observed in Georgia for first time, 312
---: Heterodera punctata, presence in Michigan confirmed, on 'Merion' Kentucky bluegrass, 399
---: lethal decline disease found in strawberry plants in Oregon, first occurrence in State, 560
---: Longidorus elongatus, (needle nematode) found in British Columbia for the first time, on strawberry, 266
---: Lophodermella cerina found on pines in Louisiana, Mississippi, Alabama and Florida for first time, 764
---: Meloidodera charis (cystoid nematode) found for first time in Nebraska, 115
---: Poria vincta var. cinerea found pathogenic to Cupressus lusitanica and Acacia melanoxylon and caused a root rot disease in East Africa, 257
---: Septoria leaf spot of pistachio, observed in Texas, not previously reported in the United States, 72
---: Septoria spot of cyclamen found in California believed to be the first occurrence in the U.S., 93
---: stem eyespot of fescues caused by Phleospora idahoensis, identified in Oregon in 1970, 63
---: stem pitting of grapevine, in Greece, 869
---: Urophlyctis leproides found on sugarbeet in California, 30
---: Xanthomonas streak (bacterial blight) of barley found in India for first time, 365
New hosts: Alternaria tenuis on ornamental herb, Barleria cristata in India, 107
---: comandra blister rust found on Pinus elliottii and P. glabra in Tennessee, 697
---: Corynebacterium michiganense on Solanum triflorum, a weed host, 336
---: wild cucumber mosaic virus, occurs in Marah oreganus, in Oregon, 546
---: Fusarium solani is cause of severe blighting of Spanish moss in the South, 390
---: Gloeosporium canker and dieback reported on Burford holly, first known report, 424
---: Helminthosporium sigmoideum found on wildrice (Zizania equatica) apparently for the first time, in Minnesota, 498
---: lespedeza cyst nematode found in yellow sweet clover, red clover, white clover (Dutch white and Ladino), and hairy vetch, 927
---: Phoma insidiosa is cause of glume blotch of wheat, 95
---: Pileolaria cotini-coggygriae is cause of smoke-tree rust on Cotinus obovatus in Arkansas, 226
---: Pythium tardicrescens found to be pathogenic to sugarcane, a first report, 766
---: new hosts of Radopholus similis in Central America, 415
New Jersey: 132; 374; 774
New Mexico: 569; 823; 1073
New York: 132; 457; 483; 565; 617; 662; 864; 874; 913
New Zealand: 45; 517; 916; 966
Nigeria: 672; 820
Nomenclature: for physiologic races of wheat stem rust, 718
Nonmercurial fungicides: for cotton seed treatments, 285; for control of Puccinia carthami on safflower seed, 1046 for barley and oats, 1122
Nonparasitic diseases: water congestion of pea, 249; 133 pea cultivars and 188 PI accessions tested for reaction, 533
North Carolina: 582; 700; 751; 956; 1013
North Dakota: 11
Nursery transplants: soybean cyst nematode controlled on transplants with nematicidal bare-root dips, 13

Oak: blackjack (Quercus marilandica), a single tree in Arkansas found with virus-like symptoms in form of chlorotic mosaic, veinbanding and chlorotic ringspots, 411; live oak, excised cankers of Endothia parasitica, treated 10 years ago in Williamsburg, Virginia have yielded a pure culture of the fungus although no canker expansion is evident, 201; species of Endothia found associated with destructive canker disease in pin oak in Virginia, 467
---: oak wilt: carried by oak bark beetles in Missouri and West Virginia after overwintering in wilt-killed trees, 108; comparison was made between the branch sampling method and the increment hammer method, for collecting samples for oak wilt, 281; caused by Ceratocystis fagacearum, trees treated with herbicides

had higher incidence and more bark beetles than untreated trees, 1106
Oats: "tulip root" of oats, caused by Ditylenchus dipsaci, trial tests in California show 'Curt' tolerant to infection but commonly grown varieties Kanota and Sierra severely injured, 433; control of covered smut of oats with carboxin, in India, 544; pathogenicity of oat crown rust isolates collected in the USA from 1966-1970, 907; several fungicides tested for control of loose smut, 1122; race 6AF was most prevalent race of Puccinia graminis avenae in the United States in 1969, 187; summary of the races of P. graminis avenae in the USA in 1970 identified by area and source of collection, 992
Ohio: 58; 333; 382; 654; 1064
Oil: chain oil of power saw, as carrier of competitive organisms to reduce the buildup of forest root pathogens, 659; commercial oils (mobil, mustard, castor and paraffin) controlled Rhizopus rot of mango fruits as a preinoculation treatment, 752; oidia of Peniophora gigantea added to SAE 30 motor oil and applied to white pine stumps resulted in better colonization in treated than untreated stumps, 834; 1.3% spray for control of pink pitting of grapefruit, 923
Oleander (Nerium oleander): previously undescribed bacterial blight of oleander caused by Pseudomonas syringae found in California in 1970 and 1971, 1030
Olive (Olea europaea): T-1 strain of Verticillium wilt appears to be associated with severe wilt in both cotton and olive in Tulare County, California, 780
Onion (Allium spp.): resistance and susceptibility of Allium species, commercial varieties and interspecific crosses to Botryotinia (Botrytis) squamosa, 394; 7-day spray interval, with maneb significantly reduced Botrytis leaf blight of onion in New York, 565; onions sprouting in storage with yellows symptoms were found to contain mycoplasma-like bodies, 824; reaction of selected onion varieties to infection by Fusarium oxysporum f. sp. cepae, 1000
Oregon: 63; 220; 263; 421; 546; 560
Overwintering: johnsongrass possibly the host in which corn stunt overwinters in South Carolina, 273; of Fusicladium dendriticum on twigs of flowering crab apple, 509; of Corynebacterium michiganense in Ohio, 654; of wheat stem rust, an unrecognized source caused uniform occurrence in late May 1970 in Kansas and Oklahoma, 990; of Helminthosporium maydis Race T in the Midwest, 563; of H. maydis in Alabama, 947; in Florida, 949; in Georgia, 951; in Mississippi, 954; in North Carolina, 956
Oxidant damage ("X" disease, "Chlorotic" decline, "ozone needle mottle"): to pines and other conifers in California showing distribution and severity of oxidant injury, in 1970, 555
Oyster plant (see Rhoeo discolor)
Ozone: low levels of ozone had little effect on growth and susceptibility of cabbage plants to Fusarium oxysporum f. sp. conglutinans, 47; distribution and severity of oxidant air pollution to native conifer species in California in 1970, 555; control of ozone concentration in an exposure chamber with a contactless meter relay, 774

Panama: 415
Pansy: Centrospora acerina found to be carried by pansy seed or in trash with the seed, 731
Papaya (Carica papaya): papaya leaf reduction virus, its transmission by Myzus persicae, 526; collar rot, a new disease of papaya caused by a Calonectria sp. with a Cylindrocladium sp. imperfect stage, 743; virus diseases in Oahu, Hawaii and identification of additional host plants, 785
Parsley: found to be a new host of alfalfa mosaic virus that develops a calico type of AMV, in California, 322; a mosaic type virus (PV) recovered from parsley and considered a host-specific strain to celery mosaic virus, 328
Pastinaca sativa (wild parsnip): western celery mosaic virus isolated from this plant in England, 268
Pathogenicity: 24 rice varieties showed wide range of reaction to some isolates of Xanthomonas oryzae, to others a narrow range, 22; pathogenic variability of Xanthomonas oryzae isolates, 593; of X. translucens f. sp. hordei to rice, 365; of isolates of oat crown rust collected in the United States, 1966-1970, 907
Pea: used in artificial system for differentiating strains of alfalfa mosaic virus, 127; damping-off and root rot, caused by Pythium aphanidermatum in Iran, thiram or captan increased emergence and yields significantly, 244; two seedling introductions PI 193586 and PI 193835 show potential source of resistant germ plasm against pea seedborne mosaic disease, 408; detection of bacterial disease in New Zealand garden pea stocks, 517; incidence of downy mildew in northern Idaho and eastern Washington, 730; Pythium ultimum was predominant species isolated

from fields in Washington with history of root rot, but was not isolated from virgin soil, 1056; water congestion of pea, in Wisconsin, 249; 533
Peach (see also Prunus): brown rot and Rhizopus rot, hot water dips or heated suspensions of benomyl or DCNA reduced infection on treated fruits, 228; Cytospora canker, development in Colorado occurs all year with the greatest enlargement in the spring, host activity rather than temperature had the greatest influence on canker enlargement, 314; D-1410 (four foliar sprays) controlled Pratylenchus penetrans in roots in New York, 617; bacterial spot, found in New Jersey and previously undescribed , is caused by Xanthomonas pruni, 774; stem pitting of peach and apricot in Greece, 1049; control of Cylindrocladium floridanum with potassium azide, 1094
Peanut (Arachis hypogaea): black rot of peanuts in Virginia, caused by Cylindrocladium spp., 419; peanut stunt virus, in Georgia, no threat to peanut production at present time, 453; leaf scorch and pepper spot, caused by Leptosphaerulina crassiasca, have distinctly different symptoms and thus are considered as separate diseases, 530
Pear: pear psylla from orchards with trees exhibiting pear decline symptoms transmitted a virus to cucumber, 295; concentric ring pattern of unknown origin found on fruit of 14 seedling pear trees at Beltsville, Maryland, 379; control of Pratylenchus penetrans in New York orchards with foliar sprays of D-1410, 617; pear latent virus, Virginia crab apple as an indicator, 983
Pearl millet: smut, caused by Tolyposporium penicillariae, effectively controlled in artificial inoculations, 87
Pecan: scab, benomyl applied at 4- and 5-week intervals effectively controlled scab in Oklahoma tests, 711
Penicillium spp.: 804; 1046
--- expansum: 791
Peniophora gigantea: 834
Pennsylvania: 509; 591; 1085
Pepper (Capsicum annuum): environmental modification of heritable resistance to potato virus Y in peppers, 604; prevalence of pepper viruses in North Carolina; 751; virus diseases of pepper in Hawaii, 783
Pepper spot: of peanut, symptom response as pepper spot or leaf scorch (caused by same pathogen) seemed to depend upon genetic makeup of plant, 530
Peppermint (Mentha piperita): black cankers and lesions on stems and rhizomes, caused by Phoma strasseri, 814
Peridermium harknessii: 163
Peronospora parasitica: 877
--- viciae: 730
Pestalotia sp.: 309; 697; 911
--- palmarum: 185
Petunia: Fusarium wilt, methyl bromide was most effective of fumigants tested, 580; control of Phytophthora in bedding plants with ETMT fungicide, 587
Pezicula malicorticis: 828
pH: fungicidal activity of potassium azide as affected by pH, 1077
Philippines: 17; 22; 216; 349; 437
Phleospora idahoensis: 63
Phloem necrosis: of elm was found in Lycoming County, Pennsylvania in 1971, 1085
Phlox divaricata: leaf spot, caused by Cercospora omphakodes, development increased after extended periods of high humidity and rainfall, benomyl gave excellent control, 1120
Phoma insidiosa: 95
--- lingam: 934
--- strasseri: 814
Phomopsis theae: 1021
Photography: a aerial photographic survey used as aid in estimating annosus root rot-caused tree mortality, 761
Phyllosticta sp.: 401; 1064
--- antirrhini: 639
Physalospora obtusa: 122
Physiologic specialization: of barley leaf rust and its sensitivity to external environment, 1026
Phytophthora sp.: 587
--- cactorum: 213
--- fragariae: 471
--- palmivora: 885
--- parasitica: 130; 929
Phytotoxicity: phytotoxicity of potassium azide (KN_3) in soil increased with increasing soil pH, 1077; of certain seed and soil fungicides potentially useful for control of cotton seedling diseases, 1111
Pierce's disease: of grape, suppression of disease symptoms with antibiotics, 610
Pileolaria cotini-coggygriae: 226
Pink disease: of cacao, caused by Corticium salmonicolor, in Solomon Islands, 885
Pink pitting: of grapefruit, caused by Cercospora sp., 923
Pinus spp.: air pollution in California, Pinus ponderosa, P. jeffreyi; P. coulteri, and P. radiata showed most serious damage over 3-year period, 555
--- banksiana (jack pine): Fomes annosus found on jack pine in Iowa, apparently for the first time, 317
--- contorta (lodgepole pine): a study of the

wave years of infection by western gall rust, 163
--- echinata (short leaf pine): estimate of tree loss from Fomes annosus, by aerial photography in Illinois, 761; spread of Fomes annosus in thinned plantations in Missouri has declined over 8-year period, particularly since 1968, 1085; decline of shortleaf pine during the past 2 years in southern Arkansas, 1036
--- elliottii (slash pine): comandra blister rust found on slash pine in Tennessee, 697; pine needle blight, first report of Lophodermella cerina on slash and loblolly pines, 764
--- glabra (spruce pine): new host of comandra blister rust in Tennessee, 697
--- jeffreyi (Jeffrey pine): Fomes annosus controlled in fresh-cut pine stumps with borax, 490
--- nigra (Austrian pine): Dothistroma pini damaged Austrian pine in Minnesota, apparently first report, 1040
--- palustris (long-leaf pine): spores of Scirrhia acicola (brown-spot needle blight) are discharged throughout year with largest numbers discharged from May to a peak in late August, 309
--- ponderosa (ponderosa pine): western gall rust, caused by Peridermium harknessii, a study of the wave years of infection in several western States, 163; borax prevented Fomes annosus colonization of fresh-cut pine stumps in California, 490; stump treatment for control of Fomes annosus, 659; Armillaria root rot, mortality decreased with increase in height of the trees, 823
--- radiata (Monterey pine): Trichoderma-treated chain oil to control Fomes annosus, 659
--- strobus (Eastern white pine): an inoculation test using Peniophora gigantea on stumps of Eastern white pine, 834
--- sylvestris (Scotch pine): brown spot needle blight in Christmas tree plantations in Wisconsin, 648; needlecast, caused by Lophodermium pinastri found in 11 States and British Columbia, 1116
--- taeda (loblolly pine): tissue from fusiform and globose shaped galls when patch-grafted onto loblolly seedlings developed galls not consistently similar to parent galls, 623; Fomes annosus losses in once-thinned plantations showed increase for about 7 years and then declined rapidly, 625; extensive needle blight in Southern States from Lophodermella cerina, first report on loblolly pine, 764; pine decline of unknown cause is affecting some loblolly pines in southern Arkansas, 1036
Pistachio: Septoria leaf spot, caused by Septoria pistaciarum, found in Texas and previously unreported in the United States, 72
Pisum sativum (see Pea)
Plasmopara halstedii: 11
Ploioderma hedgcockii: 764
--- (Hypoderma) lethali: 764
Podosphaera leucotricha: 514
Poinsettia: control of Thielaviopsis basicola root rot with benomyl and thiabendazole, 238
Poison hemlock (Conium maculatum): strain of celery mosaic virus found in poison hemlock in California, 328
Poland: 445; 828; 932
Populus deltoides (eastern cottonwood): Septoria canker on nursery stock in Mississippi, 460
Populus tremuloides (quaking aspen): aerially applied antibiotics fail to control Hypoxylon canker in Minnesota, 650; Hypoxylon canker, Fomes igniarius heart rot, and Nectria canker found to be three major diseases in Adirondack region of New York, 662
Poria vincta var. cinerea: 257
Postharvest decay: of apples: storage decay from Nectria galligena controlled with 2 AB (2-aminobutane); thiabendazole, captan and DCNA not effective, 771; of mechanically picked apples, controlled with thiabendazole, 791
---: of cabbage, Botrytis cinerea and Sclerotinia sclerotiorum were apparently controlled in a 1.4% oxygen storage atmosphere; Fusarium roseum was still active, 864
---: stem-scar and rind mold of cantaloup controlled with fungicides better than with a hot water treatment, 183
---: of citrus, decay of degreened oranges and tangerines as affected by time of washing and fungicide application, 149
---: of mahgo, screening trials for control of Diplodia and Rhizopus rots, 752
---: of nectarine: heated and unheated benomyl dips equally effective in control of brown rot if fruit had preharvest spray, otherwise, the heated dip was significantly better, 69; reduced with preharvest benomyl sprays and postharvest wax-fungicide treatments, 804
---: of peach, control of brown rot and Rhizopus rot of inoculated peaches with hot water or hot chemical suspensions, 228
---: potato, fungicidal dips or dust reduced tuber dry rot of potato caused by Fusarium roseum sambucinum, 723
---: of strawberry, control of Botrytis rot on

fruit with benomyl dip, 883; strawberry nursery plants treated with benomyl resulted in 85% control of plant decay after 7 months' storage, 1009
Potato: seed and soil treatments compared for control of Rhizoctonia solani on stems and stolons, 550; tuber dry rot, caused by Fusarium roseum sambucinum, reduced with fungicides, 723; Verticillium albo-atrum infection of potatoes was delayed by fumigation of soil with 1,3-D + chloropicrin in Connecticut, 968
Potato virus X and Y: on pepper in North Carolina, 751
Potato Y virus: on pepper, 783
Powdery mildew: of apple: control with fungicides as influenced by seasonal temperatures, 263; benomyl and triarimol appear to be adequate substitutes for Karathane in control of apple powdery mildew, in Washington, 514
---: of black currant, controlled in Poland with benomyl, 932
Practical control: of lawn grass diseases by removal of mower clippings, 778
Pratylenchus penetrans: 276; 617; 673; 968; 1032
--- zeae: 41
Precipitation: relationship of temperature and precipitation to the distribution of Claviceps gigantea on corn in Mexico, 231
Preharvest decay: of nectarine, two bloom sprays of benomyl reduced brown rot infection, but did not protect fruit against postharvest decays, 69; preharvest spray to control decay of nectarines both before and after harvest, with benomyl, 804
Preharvest infection: of cotton fiber, distribution of microbial infections in the U.S. Cotton Belt, 714
Preharvest treatments: of strawberry plants controlled storage decay, 1009
Premature foliation: of peach and apricot, affected with stem pitting, in Greece, 1049
Primrose: Ramularia leaf spot controlled on hybrid primrose with benomyl, 586
Prunus (see also common names): bacterial canker on orchard trees in Oregon, the susceptibility of the root or trunk stock is related to the severity of the disease of the scion cultivar, 421; indexing rose mosaic in Rosa indica on 'Shirofugen' flowering cherry for certified planting stock is apparently of limited reliability, 431; 12 suspected viruses were not eliminated by hot water treatment of budsticks, 738
Pseudomonas sp.: 361; 447; 517
--- lachrymans: 3
--- marginalis: 1037
--- mors-prunorum: 961
--- phaseolicola: 54; 203; 703
--- solanacearum: 521
--- syringae: 421; 703; 961; 1030
Pseudopeziza ribis: 932
Pseudopityophthorus minutissimus: 108
Pseudotsuga macrocarpa: air pollution damage, in California, 555
Psylla pyricola: 295
Puccinia sp.: 317
--- carthami: 1046
--- coronata var. avenae: 907
--- graminis f. sp. avenae: 187; 992
--- --- f. sp. tritici: 187; 695; 871; 986; 990
--- hordei: 1026
--- purpurea: 941
--- recondita: 172
--- --- tritici: 691; 718
--- zoysiae: 778
Puerto Rico: 506; 902
Pyricularia grisea: 778
Pythium sp.: 587
--- spp.: 536; 837; three unidentified Pythium species were isolated from virgin soils collected in eastern Washington; none was pathogenic to beans or peas, 1056
--- aphanidermatum: 244; 851
--- debaryanum: 851
--- graminicola: 766
--- tardicrescens: 766
--- ultimum: 851; 1056

Quercus (see Oak)

Races: of downy mildew of lettuce, over 50 lettuce plant introductions were immune to races 5 and 6 of Bremia lactucae, 794
---: of Helminthosporium maydis: sclerotia formation by H. maydis Race T on corn seed, 755; Race T found in Nigeria, West Africa, 672; distribution of Race T, 302; use of toxic filtrate from Race T cultures to test for resistance in corn seedlings, 469; overwintering of Race T in Midwestern United States, 563; 934 isolates of H. maydis were identified as to race, relative virulence and mating type, in the Northeastern United States, 495; Races O and T have been isolated and identified in Hawaii on basis of reaction on differential corn lines, 392; in Iowa in 1968, in stored ear corn, 855
---: of Hemileia vastatrix, race II of coffee rust is the race present in Brazil in contrast to races I, II, VII, and XV, on Coffea arabica, and IV on C. canephora, in Kenya, 209
---: of Puccinia coronata avenae (oat crown rust) in the U.S. from 1966-1970, 907

---: Puccinia graminis f. sp. avenae, Race 31 comprised 82% of 1391 isolates collected in the USA in 1970, 992
---: Puccinia graminis f. sp. tritici: races 15B-2 and 11 accounted for 55 and 15%, respectively, of isolates identified from uredial collections in 1969 in the United States, 187; Race 15, 15B-2, 15B-6 and 151 were the most prevalent races in the USA during 1970, 986
---: Puccinia hordei, Races 4, 40, 44, and 53 identified in Minnesota, 1026
---: Puccinia recondita tritici, system of nomenclature for physiologic races, 718
---: of burrowing nematode, Radopholus similis in Central America, 415
---: new race of Uromyces phaseoli var. vignae (cowpea rust) found in south Georgia, 384
Radopholus similis: 154; 415
Rain: injury to tissues of upper pulvini of petioles of primary leaves of bean resulted from heavy rain and caused severe defoliation to a commercial dry bean planting in New York, 457
Ramularia carthami: 846
--- primulae: 586
Raspberry: fruit rot, Cuthbert, Meeker and Matsqui cultivars showed least susceptibility to fruit rot of 25 cultivars tested, 135; sodium salt of dehydroacetic acid (DHA-S) sprayed on fruit before harvest extended the shelf-life of red raspberries by several days, 175
Redbud (Cercis canadensis): redbud planted in soil heat-treated prior to infestation with Verticillium albo-atrum developed greater incidence of disease than in unsterilized soil, 996
Relative humidity: combination of high moisture, RH, and warm temperatures contribute to occurrence of water congestion of pea in Wisconsin, 249
Reproduction: of sugarbeet nematode and cabbage root nematode on several cultivars of Brussels sprouts, 1005
Resistance: apple, to Nectria canker, varieties Beauty of Rept, Boiken, Close and Wealthy of 33 varieties tested in Poland were free from cankers under natural conditions, 445; barley cultivar Marnobarb was immune to Helminthosporium stripe in greenhouse tests and six other cultivars showed less than 1% infection, 858; corn, spray and injection techniques for rapid screening of large numbers of plants for resistance to Helminthosporium maydis, 469; cotton, resistance in commercial varieties to Fusarium wilt, 896; cucumber, several pickling type varieties showed varied levels of resistance to Corynespora leaf spot, none found in slicing types, 142; pea, screening for resistance to pea seedborne mosaic virus, 408; breakdown of resistance to Fusarium wilt in 'Wisconsin Perfection' peas in presence of Pratylenchus penetrans, 1032; pepper, modification of heritable resistance to potato virus Y in peppers, 604; in Krish sorghum to sugarcane mosaic virus, 596; strawberry, a method of screening strawberry seedlings for resistance to Phytophthora fragariae, 471; resistance in strawberry varieties and lines to Mycosphaerella fragariae in Illinois, 573; tomato, inheritance of tomato anthracnose resistance, 253; Ulmus pumila and U. parvifolia showed highest resistance to Dutch elm disease of nine elm species evaluated, 980; 35 cultivated crops tested for resistance to Meloidogyne incognita and M. javanica, 1082
Rhizoctonia sp.: 614
--- fragariae: 599
--- solani: 347; 550; 861; 864; 1077; 1111
Rhizopus sp.: 1097
--- arrhizus: 752
--- stolonifer: 69; 228; 714; 804
Rhoeo discolor (oyster plant): tan leaf spot, caused by Curvularia eragrostidis, in Florida, 38
Rhopalosiphum maidis: 874
--- padi: 874
Rice: Curvularia cymbopogonis was found on rice seed samples from the Philippines, previously unreported there, and a first record of its pathogenicity to rice, 349
---: bacterial diseases: varietal resistance to bacterial blight, caused by Xanthomonas oryzae, 17; pathogenicity of bacterial blight pathogen of barley (Xanthomonas translucens f. sp. hordei) to rice and its possible close relationship to the rice streak pathogen, X. translucens f. sp. oryzae (X. oryzicola), 365; inoculation of rice with the pathogen of bacterial leaf streak of rice by sandpaper and sand rubber plate method, 404; pathogenic variability among Xanthomonas isolates indicates some host specialization in pathogen and therefore the advisability for use of several isolates in screening rice varieties for resistance, 593
---: stem rot, caused by Sclerotium oryzae, occurs throughout the northern California rice-growing regions, 757
Rind mold: of cantaloup, control with fungicides, 183
Root dips: citrus seedlings treated by bare-root dips with several nematicides were protected against burrowing and citrus nematodes, 154

Rose (Rosa multiflora japonica): control of root-knot nematode, Meloidogyne hapla, on 2-year-old plants with D-1410, 89; rose spring dwarf is a graft-transmissible disease that is being increased through use of both infected sionwood and infected rootstock cuttings, 294; use of 'Shirofugen' flowering cherry as reliable indicator for rose mosaic virus is limited, thus a larger number of indexings needed for each rose variety, 431; comparison of symptoms induced by rose wilt virus and Verticillium dahliae, 916
Rot: bacterial soft rot, of bok choy, caused by Pseudomonas marginalis, 1037
---: basal rot: of Iris bulbs, 425; Fusarium basal rot of Narcissus, effective controls with benomyl and thiabendazole, 428; Fusarium basal rot of onion, four commercial varieties, tested together for time of initial root infection and level of infection, reacted similarly, 1000
---: basal sprout rot, of Brussels sprouts, disease apparently associated with maturity, 297
--- black rot of sugarcane cuttings in Hawaii, 7
---: black, of peanut, caused by Cylindrocladium spp. in Virginia, 419
---: black rot of sweetpotato, thiabendazole was most promising fungicide for control of Ceratocystis fimbriata, 523
---: boll rot: cotton: distribution of Aspergillus flavus boll rot and its association with fluorescence of the fiber, 33; estimate of crop loss in 1970 from boll rots, 504; distribution of Aspergillus flavus boll rot in the U.S. cotton crop for 1970, 510
---: brown rot of nectarine, pre- and postharvest control with benomyl, 69; of peach, 228
---: collar, of papaya, a new disease, 743
---: corm rot of gladiolus, latent infection by Fusarium activated effectively by CO_2 treatment, 340
---: crown, of apple, caused by Phytophthora cactorum, 213
---: ear, and stalk rot of corn, 158
---: fruit: apple, source and longevity of inoculum of Botryosphaeria ribis and Physalospora obtusa, 122; raspberry, susceptibility of red raspberry cultivars to fruit rot, 135; of mango, Diplodia stem-end rot controlled with thiabendazole, 752; of mango fruits, Rhizopus controlled with commercial oils in preinoculation treatment, 752; of peach, Rhizopus rot controlled with heated chemical suspensions and heated water, 228
---: kernel and ear rots of corn, are more severe this year with the production of high-lysine corn that is very susceptible ear rot pathogens, 1046
---: root: Fomes annosus: in Iowa, 317; stump treatment for prevention, 490; mortality in once-thinned loblolly pine, 625; stump treatment for biological control, 659; aerial photographic survey for loss, 761; loss in thinned shortleaf pine, 1085
---: ---: Armillaria root rot in four ponderosa pine plantations in New Mexico, 823; Chinese gooseberry vines susceptible to Armillaria mellea at U.S. Plant Introduction Station, Chico, California, 1099
---: ---: bean root rot, caused mainly by Fusarium solani f. phaseoli, 483
---: ---: black root rot of bean and tobacco, caused by Thielaviopsis basicola, 352
---: ---: Cylindrocladium root rot of seedling tulip-poplar in North Carolina nursery, 700
---: ---: foot and root, of citrus, caused by Phytophthora parasitica, 929
---: ---: of forest trees, Cupressus lusitanica and Acacia melanoxylon in East Africa, 257
---: ---: of pea, caused by Pythium aphanidermatum, in Iran, 244
---: ---: Phymatotrichum omnivorum, loss estmiate of cotton crop for 1970, 504
---: ---: of Poinsettia, caused by Thielaviopsis basicola, 238
---: ---: Pythium: Pythium brown rot of carrots in muck soil in Wisconsin, 536; Pythium root rot of chrysanthemum, 851; of sugarcane, individual and combined effects of Pythium tardicrescens and P. graminicola on sugarcane, a first report, 766; of pea and bean, Pythium ultimum apparently develops rapidly in land cropped to a susceptibile host, 1056
---: ---: Sclerotium root rot, of sugarbeet, 630
---: stalk: Diplodia ear and stalk rot of corn are apparently independent characters in a general sense, based on correlation differences in five studies, 158; of corn, 634
---: stem: of wildrice (Zizania aquatica), possibly first report of Helminthosporium sigmoideum on Z. aquatica, 498; of rice, caused by Sclerotium oryzae, distribution and occurrence in California, 757; basal stem rot, of hardwoods in West Pakistan, caused by Ganoderma lucidum, 1130; rhizome and stem rot of peppermint, caused by Phoma strasseri, 814
---: tuber dry rot, of potato, reduced with fungicidal dips or as dust, 723
Rotylenchulus reniformis: 362; 1043
Rubber tree (Hevea brasiliensis): bark necrosis, caused by Fusarium solani and Bo-

tryodiplodia theobromae in Malaysia; can be controlled with benomyl, captafol, and an organo-mercury fungicide, 152
Rust: a proposed system for coding infection types of the cereal rusts, 319
---: of black currant, Cronartium ribicola controlled with benomyl, 932
---: blister, Cronartium comandrae infection found on slash pine and spruce pine in Tennessee, apparently a new record, 697
---: coffee: caused by Hemileia vastatrix, in Brazil and Kenya, 209; control of coffee rust in Philippines, 216
---: of cowpea, an apparent new race of cowpea rust, Uromyces phaseoli var. vignae, was isolated from southern peas in south Georgia, 384
---: crown rust of oat, race group 264B was most common race group in the USA in 1970, 907
---: gall: wave years of infection of western gall rust on pine, 163; rust galls on loblolly pine, fusiform and globose-shaped galls grafted onto seedlings produced variations in gall shapes that indicate that the pathogen alone does not determine the gall shape, 623
---: leaf: of barley, variation in rust sporulation due to external environment, and effect of maleic hydrazide indicate several infection type readings are advisable for noting infection types, 1026; of sorghum, from 496 selections from the Rockefeller-India collection, 12 were resistant and 9 were highly resistant, 941; of wheat, the distribution of virulence in the wheat leaf rust population in North Dakota in 1970, 172; of wheat, Puccinia recondita f. sp. tritici controlled in growth chamber study with new systemic fungicide, Cela W-524, 961; nomenclature of important races of P. recondita tritici will be based on a "type" culture permanently preserved in liquid nitrogen, 718
---: Pileolaria cotini-coggygriae on smoketree (Cotinus obovatus), in Arkansas, 226
---: stem: of oats, 187; of oat, races of Puccinia graminis f. sp. avenae in the United States in 1970, 992; of wheat, identification of new physiologic races of P. graminis f. sp. tritici in the United States in 1969 and their prevalence and distribution, 187; races of P. graminis f. sp. tritici of the United States in 1970, 986; evidence for an unrecognized source of overwintering of stem rust of wheat in the U.S., 990
---: Zoysia, control on lawn grass by prompt removal of grass clippings in Mississippi, 778
Rye: nonhost for Meloidogyne incognita, used as winter cover crop before muskmelon crop in Indiana apparently reduces need for fumigation, 55

Safflower: Cercospora carthami and Ramularia carthami, leaf spot pathogens produce conidia that can withstand prolonged desiccation, 846; maneb and TCMTB controlled seedborne safflower rust and appear suitable substitutes for alkyl mercury compounds, 1046
Sainfoin (Onobrychis viciaefolia): sainfoin (possible replacement crop for alfalfa) was similar to alfalfa in susceptibility to northern root-knot nematode in Utah, 1069
Samoa: 50
Scab: apple: imperfect stage, Fusicladium dendriticum, on twigs of flowering crab apple in Pennsylvania, 509; benomyl, NF-44 and NF-35 were comparable to dodine, in control of apple scab, 828
---: sour orange scab, benomyl gave better control than copper spray on citrus hybrid 'Page', in Florida, 251
---: pecan, Fusicladium effusum effectively controlled with benomyl, 711
Scaphoideus luteolus: 1085
Scaphytopius acutus: 645
Schizaphis graminum: 874
Scirrhia acicola: 309; 648
Sclerospora sorghi: 202
Sclerotia formation: by Helminthosporium maydis Race T, 755
Sclerotinia spp.: 1087
--- oryzae: 757
--- rolfsii: 630; 861
--- sclerotiorum: 234; 864; 1077; 1136
Scolytus multistriatus: 271
Scotch tape: for the trapping and examination of spores, 832
Scurf: of sweetpotato, water and ferbam treatments at 55°C for 1/2 min or at 49° for 5 min controlled scurf, 746
Seed: Centrospora acerina carried by pansy seed, 731; germination tests of corn seed lots grown in Minnesota in 1970 which contained Texas male-sterile cytoplasm averaged 10% kernels infected with Helminthosporium maydis, 371; germination tests on commercial samples of corn grown in 1970 detect infection with H. maydis, 726; sclerotial formation of H. maydis Race T on corn after 5 days in germination tests, 755
---: transmission, possible seed transmission of H. maydis and Fusarium moniliforme by infected corn seed, especially that with T cytoplasm, 486
---: treatment: anaerobic and hot water treatments reduced germination of barley

seeds, within 12 months of treatment, 118; several nonmercurial fungicides are effective as cottonseed protectants to replace canceled alkyl mercury compounds, 285; for control of bean root rots, 483; corn seed treated with carboxin reduced head smut, 501; for control of some important cereal smuts in India, 544; seedpieces of potato treated with PCNB and benomyl reduced infection from Rhizoctonia solani, 550; 15 chemicals controlled loose smut of oats on 'Early Ballard'; only potassium azide and sodium azide controlled loose smut of barley when applied to naturally infected seed; several fungicides controlled barley leaf stripe 1122

Seedborne: seed infection of popcorn with Helminthosporium maydis, 382; seed infection of corn from Indiana by H. maydis, 387; H. maydis introduced on seed corn is suggested for the occurrence of H. maydis in widely scattered and isolated plantings in Hawaii, 392; interception of Race T of H. maydis on imported corn seed lines and gift packs of popcorn from the United States in New Zealand, 966; seedborne fungi, Alternaria tenuis, Phyllosticta antirrhini, and Stemphylium botryosum found to be pathogenic to Antirrhinum majus, 639; first report of Thielavia basicola and Pestalotia sp. as seedborne pathogens of soybeans in the U.S., 911

Seedling: defoliation of bean caused by heavy rain, 457; alfalfa, normal stem nematode population development in inoculated seedlings may be reduced in an area being treated with dichlorvos pest strips, 621; cotton species and varieties tested for susceptibility to Colletotrichum gossypii; 879; American elm, adjuvants added to benomyl soil drench did not increase uptake of chemical in plants grown in planting media, 971

Seedling diseases: seedling injection as an aid in identifying bean blight bacteria, 703

---: of corn, Helminthosporium maydis seedling blight of popcorn, 382

---: of cotton: fungicidal sprays applied over the top of cotton rows compared with "wet" and no treatment "dry" controls showed that a valid comparison of aqueous fungicidal sprays and "dry" planting cannot be made, 307; damping off, of cotton, estimated loss by State, 504; of cotton, in vitro method of comparing systemic fungicidal activity and phytotoxicity of fungicides potentially useful for disease control, 1111

---: Pinus palustris infection by Scirrhia acicola spores greatest in Mississippi from May to August with a peak of ascospore production in late August, 309

Septoria cyclaminis: cause of Septoria spot of cyclamen, not previously reported from the United States, 93

--- musiva: 460

--- pistaciarum: pistachio nut trees in Texas found susceptible to this pathogen in the form of severe leaf spotting, a new disease to the United States, 72

Sigatoka (see Banana leaf spot)

Smoke-tree (Cotinus obovatus): rust, caused by Pileolaria cotini-coggygriae, found on native smoke-tree in Arkansas, a new host species, 226

Smut: covered smut of oat and barley, caused by Ustilago hordei, in India, seed treatments with carboxin gave almost complete control, 544

---: head: of sorghum: pathogen, Sphacelotheca reiliana, observed at Blairsville, Georgia in September 1970, for first time, 312; of corn, reduced with carboxin seed treatments, 501

---: loose smut of wheat, controlled with benomyl and carboxin, 1053

---: of pearl millet in India, oxycarboxin was better than other fungicides tested for control of Tolyposporium penicillariae, 87

Snapdragon (see Antirrhinum)

Soil: amendments: alfalfa hay, leaves and stems of cabbage, castor pomace, corn stover, and chestnut tannin at .5% of air-dry weight of soil reduced black root rot of tobacco, 352; mustard, Brussels sprouts, kale, collards, cress, cabbage, turnip, kohlrabi, rhubarb, Swiss chard and soybean dry tops and stems added to greenhouse soil reduced both pre- and post-emergence damping-off of sugarbeets, 440; green crop amended soil caused root damage to lettuce seedlings in field and greenhouse tests in California, 837; oat straw and mineral fertilizer reduced severity of root knot of tomato, caused by Meloidogyne incognita, 1126

---: treatments: effect of adjuvants on uptake of benomyl from planting media by American elm seedlings, 971; with potassium azide, relation of soil pH to effectiveness, 1077

---: ---: for fungi: benomyl ineffective against Dutch elm disease, by soil application, 58; soil injection of benomyl against Dutch elm disease, 179; 11 fungicides tested for possible use in field trials to prevent crown rot of apple, caused by Phytophthora cactorum, 213; benomyl and thiabendazole added to soil controlled root

rot of Poinsettia, caused by Thielaviopsis basicola, 238; benomyl and thiophanate as soil drenches controlled Fusarium blight on bluegrass turf, 333; benomyl + chloroneb in-furrow sprays were more effective than seed treatments in control of bean root rots, 483; benomyl, chloroneb and PCNB as soil treatments were more effective than seed treatments for control of Rhizoctonia infection on stems and stolons of potatoes, 550; field treatments with benomyl + surfactants to control Verticillium wilt of cotton, 569; soil fumigation for control of Fusarium wilt of petunia, 580; ETMT fungicide as a soil drench or soil mix was superior to Dexon in control of seedling diseases of bedding plants, caused by Pythium or Phytophthora, 587; soil drenching with tetracycline suppressed disease symptoms of Pierce's disease in grape, 610; ridge-soil drenching with carboxin and chloroneb reduced Sclerotium root rot of sugarbeet, 630; cutting drench treatment with benomyl controlled Cylindrocladium rot of azalea, 679; for control of Phytophthora parasitica in citrus orchards, 929; Verticillium infection of barberry and redbud was increased when soil was heat-treated prior to infestation with pathogen, 996; benomyl added to soil of potted tea plants was readily taken up by plants and translocated, 1021; with potassium azide for control of Cylindrocladium floridanum, 1094

---: ---: for nematodes: fumigation for control of root knot before planting muskmelons, 55; drench treatments of D-1410 apparently eradicated Meloidogyne hapla from field-grown roses, Rosa multiflora japonica, 89; aldicarb and BAY 68138 effectively controlled lesion nematodes in heavy greenhouse soils, 276; DBCP drenches for control of nematodes in banana in Cyprus, 707; nematicide fumigation of cabbage fields on nematode infested soil increased yields, 841; fumigation with 1,3-D or 1,3-D + chloropicrin controlled Pratylenchus penetrans nematodes and suppressed infection from Verticillium albo-atrum on potato resulting in increased yields, in Connecticut, 968; for control of Heterodera schachtii, nematicides 1,3-D and DD effective but depth applied was not significant, 1101

Solanum triflorum: a common solanaceous weed in southeastern Wyoming and growing in tomato breeding plots was found to be naturally infected with bacterium Corynebacterium michiganense, 336

Solomon Islands (Brit.): 885

Sorghum: Krish sorghum (S. halepense x S. roxburghii) showed high resistance to four strains of sugarcane mosaic virus in Australia, and maybe a possible source of resistance for grain and fodder sorghums, 596

Sorghum (Sorghum bicolor): head smut of sorghum (Sphacelotheca reiliana) found for first time in Georgia, 312; head mold, in 1969 and 1970 head mold was primarily incited by Fusarium spp. in Kentucky, 337; sources of resistance in sorghum to Puccinia purpurea, the causal agent of leaf rust, 941; a mechanically transmissible virus isolated from sorghum in southern France, 1090

South Africa: 747

South Carolina: 158; 273; 741; 810; 1122

South Dakota: 172

Southern corn leaf blight (SCLB): survival and overwintering properties of the pathogen, Helminthosporium maydis, 99; yield reductions, from plants with various cytoplasm content, 115; corn inbreds with different cytoplasms varied in susceptibility to H. maydis with the type of cytoplasm; lines with T cytoplasm were susceptible and lines with normal or J cytoplasm were resistant, 195; geographic distribution of H. maydis Race T for 1969 and 1970, 302; development of seedling wilt of corn in kernels infected with H. maydis in Minnesota, 371; H. maydis found to be seedborne on popcorn seed from blight-affected planting in Ohio, 382 seed samples from the 1970 corn crop from Indiana indicate significant amounts of SCLB infection, 387; in Hawaii, the spread since first record of occurrence in 1968 suggests independent introductions in seed corn, 392; method for testing large numbers of corn plants for resistance to blight pathogen, 469; the effect of cytoplasmic inheritance on susceptibility of corn hybrids and the importance of diversity of cytoplasms in a breeding program, 473; severe pathogenicity of H. maydis to corn seedlings, especially those with T cytoplasm, 486; identification of 934 isolates as to race frequency, distribution and virulence in 1970, 495; overwintering of Race T in 1970-71, in South Dakota, Minnesota, Wisconsin, Michigan, Ohio, Indiana, Illinois, Kansas, Nebraska and Missouri, 563; H. maydis T-toxin as an indicator of resistance to SCLB, 667; a severe attack of leaf blight occurred in Nigeria in 1970 on widely-grown cultivar, NS-5, 672; detection of kernels infected with H. maydis in commercial

samples of corn produced in 1970, 726; sclerotial formations by H. maydis Race T on corn seed, 755; H. maydis isolates collected in 1970 from Southeastern U.S. were genetically more diverse than isolates from the Midwest, 759; Race T was discovered in commercial corn produced in 1968 in Iowa, 855; overwintering of H. maydis in Alabama, 947; in Florida, 949; in Georgia, 951; in Mississippi, 954; in North Carolina, 956; corn seed from the United States infected with blight pathogen intercepted in New Zealand, 966
Southern pea (see Cowpea)
Soybean: association of pod and stem blight with stem breakage, 776
---: nematodes: apparently a nonhost of lespedeza cyst nematode, 974; breeding lines resistant to reniform nematode are also resistant to soybean cyst nematode but all lines resistant to soybean cyst nematode are not resistant to reniform nematode, 1043
---: Phytophthora parasitica, six varieties inoculated with pathogenic isolate of P. parasitica from white clover show differences in susceptibility, 130
---: Thielavia basicola and Pestalotia sp. found internally seedborne in soybean, not previously reported from the U.S. in this crop, 911
---: virus, bean pod mottle virus was successfully transmitted to soybean by striped blister beetle in greenhouse tests, first record of this species as vector of BPMV, 628
Spanish moss (Tillandsia usneoides (= Dendropogon usneoides)): a stem and leaf blight caused by Fusarium solani threatens the existence of Spanish moss in some regions of the Southeastern United States, 390
Sphaceloma fawcetti: 241
Sphacelotheca reiliana: 312; 501
Sphaerotheca lanestris: 342
Spore dispersal: of Scirrhia acicola in Mississippi, ascospores were discharged during rains, dews and fogs, with low production at low temperatures and an increase from early spring to a peak in late August, 309; use of Scotch tape as the adhesive-coated trapping substrate on a rotating drum spore trap, 832
Spotted wilt virus: of pepper, 783
Sprinkler irrigation: Cercospora leafspot of sugarbeet controlled best with benomyl under sprinkler irrigation, 449
Statice (Limonium sinuatum): falling water drops appear to be the direct causal factor of flower blast of statice, 848
Stem breakage: of soybean, branches and pods in contact with soil were much more severely infected with Diaporthe phaseolorum var. sojae than erect stems, 776
Stem-scar: of cantaloup, benomyl gave better control than other fungicides tested, or a hot water treatment, 183
Stem dieback: of yellow birch seedlings in Wisconsin and Michigan, 788
Stem eyespot: on native fescue, caused by Phleospora idahoensis, found for the first time in Oregon, 63
Stem pitting: of grapevine, in Greece, 869; of peach and apricot in Greece, 1049
Stemphylium botryosum: 374; 639
St. Augustine blast or gray spot: Pyricularia grisea controlled on lawn grass by removal of grass clippings, 778
Storage: barley seeds treated anaerobically and with hot water showed reduction in seedling emergence within 12 months from time of treatment, 118; genetic stability of Puccinia graminis tritici in cryogenic storage, 695; of onions, sprouting onions in storage contained mycoplasma-like bodies, 824; controlled-atmosphere storage for cabbage, 864; 10 years of storage of urediospores of Puccinia graminis f. sp. tritici in liquid nitrogen has caused little deterioration in viability, 871; of conidia of apple scab fungus by freezing, 976
Storage decay (see Postharvest decay)
Strains: of alfalfa mosaic virus, an artificial system for differentiation, 127; presence of watermelon mosaic viruses 1 and 2 in squash in New York and New Jersey is confirmed, 132; ash strain of tobacco ringspot virus transmitted by nematodes, 475; new strains of Xanthomonas phaseoli (common blight of bean) from Colombia reported, 505; of sugarcane mosaic virus, Krish sorghum shows high resistance to four strains, 596; Verticillium albo-atrum isolates from Australia compared with SS-4 and T-1 strains from cotton grown in South, Central and North America, 977
Strawberry; first report of natural occurrence of lethal decline disease on strawberry outside of Washington State, in Oregon, 560; control of storage decay of strawberry plants with benomyl preharvest treatment, 1009
---: angular leaf spot, caused by Xanthomonas fragariae, identified in Florida, 142
---: anthracnose, caused by Colletotrichum fragariae, controlled with benomyl + pinolene, 139
---: Botrytis cinerea, benomyl controlled Botrytis rot of strawberry fruits if fruit was dipped 24 hours prior to or at time of artificial inoculation, not when dipped

24 hours later, 883
---: leaf spot, resistance of strawberry varieties and selections to Mycosphaerella fragariae, 573
---: needle nematode, discovered in Fraser Valley of British Columbia in field of strawberries in fall of 1969, 266
---: Phytophthora fragariae: liquid zoospore cultures utilized for screening seedlings for resistance, 471
---: Rhizoctonia fragariae, a mycorrhizal and pathogenic fungus of strawberry plants, 599
---: virus, study of virus level and horticultural quality of strawberry runner plants grown in Oregon for the 2nd year of a 3-year study, 220
Strelitzia reginae: bacterial leaf stripe in Florida, caused by Pseudomonas sp. differs from bacterial wilt of Strelitzia in Hawaii, caused by P. solanacearum, 447
Stump treatment: of ponderosa pine and Jeffrey pine stumps with borax to prevent Fomes annosus infection, 490; Trichoderma sp. applied through chain oil of power saw to control Fomes annosus in pine stumps not successful, 659
Sugarbeet: Urophlyctis leproides, caused tumors on leaves and crowns of beet in California, 30
---: Cercospora beticola: control of Cercospora leaf spot by aerial spraying with benomyl, in Israel, 408; control of leafspot under sprinkler irrigation with benomyl, 449; effective fungicide control of C. beticola in Iowa and Nebraska with timed sprays, 479; epidemics on sugarbeets in the Texas-New Mexico High Plains, 1073
---: damping-off, caused by Aphanomyces cochlioides, reduced with various fungicides and soil amendments, 440
---: nematodes, 1,3-D and DD controlled Heterodera schachtii, depth of application showed no significant effect, 1101
---: root rot, control of Sclerotium root rot with systemic fungicides, 630
---: virus: curly-top virus, found in Iran in 1966 is apparently spreading through the sugarbeet-growing areas of the country, 85; incidence and distribution of aphid-borne viruses of sugarbeets in Washington, 103; relationship between volunteer sugarbeets and occurrence of beet mosaic and beet western yellows virus in Washington beet fields, 676
Sugarcane: black rot of sugarcane, Ceratocystis adiposa, was isolated for the first time in Hawaii, 7; pathogenicity of Pythium tardicrescens to sugarcane demonstrated for first time in greenhouse tests, 766
Sunflower (Helianthus annuus): a serious outbreak of downy mildew, caused by Plasmopara halstedii, in the principal sunflower production area in the U.S., 11; found to be host of Leveillula taurica, in Iraq, 192; first report of Fusarium wilt of sunflower in southwest Texas, 1124
Survival (see also Viability): Helminthosporium maydis isolates: ability to survive in parasitic and overwintering phases, 99; winter survival of H. maydis in five States, 947; 949; 951; 954; 956
---: of apple fruit rot inoculum, Botryosphaeria ribis and Physalospora obtusa under orchard conditions in Virginia, 122
Susceptibility: of red raspberry cultivars to fruit rot, 135; of Cucurbita spp. to root-knot nematode, Meloidogyne incognita, 666; of Gossypium species and varieties to seedling anthracnose, 879; hypersusceptibility of high-lysine corn to kernel and ear rots, 1046; of common sainfoin to Meloidogyne hapla, 1069
Sweetpotato: black rot, caused by Ceratocystis fimbriata, thiabendazole showed good control of fungus, 523
---: scurf, water and ferbam dips at 55°C for 1/2 min and at 49° for 5 min controlled scurf, 746
---: Rhizopus rot, DCNA dip effective in control of soft rot in root pieces of 'Redmar' and 'Julian' cultivars, 1097
---: virus, sweetpotato virus infections in Puerto Rico, 902
Swordfern (Nephrolepis exaltata 'Florida Ruffle'): a blight of fronds found to be caused by Rhizoctonia sp., 614
Sycamore: (Platanus occidentalis): anthracnose epidemic in Mississippi-Yazoo Delta occurred during the spring of 1970, 93; Diplodia theobromae and Ceratocystis fimbriata f. platani found in silage sycamore plantings in Georgia and South Carolina, 741
Symptomatology: of coffee rust in Brazil and Kenya, 209; symptomatology, host range studies, electron microscopy, and serology tests show wild cucumber mosaic virus infects Marah oreganus, 546

Tea plant (Thea sinensis): uptake of benomyl by tea plant and its effectiveness against stem canker, caused by Phomopsis theae, 1021
Temperature: effect of dip temperatures on symptom expression of dip-inoculated tomato seedlings with tobacco mosaic virus, 375; dip or soak temperatures of benomyl and thiabendazole raised from

10 to 40°C improved control of Fusarium basal rot of Iris bulbs, 425; heat treatments to eliminate virus symptoms in sweetpotato clones in Puerto Rico failed, 902
---: air: mean annual temperatures of 13-15° C and total annual precipitation of more than 1000 mm favor development of Claviceps gigantea on corn in Mexico, 231; high temperature and excessive soil moisture in Iran contribute to development of serious damping-off and root rot of pea, 244; warm temperature, coupled with high soil moisture and high RH apparently favor the nonparasitic disease of pea, water congestion, 249; cold pox of butternut squash observed in Florida in 1970, severity increased with decreasing temperature, 350; relation of temperature to development of Cercospora leafspot epidemics, 1073
---: hot water treatment: hot water + benzimidazoles and ethephon produced larger and heavier gladiolus corms, 82; and anaerobic treatments reduce longevity of barley seeds, 1118; effect of hot water and fungicides for control of stem-scar and rind mold of cantaloup, 183; and benomyl or DCNA heated dips significantly improved decay control over unheated water or chemical dips, for control of brown rot or Rhizopus rot of peach, 228; soaking Prunus budsticks at 50°C for 15-20 min did not eliminate 12 suspected viruses, 738; sweetpotatoes dipped for 1/2 min at 55°C for 5 min at 49° gave good control of scurf, 746
---: soil, heat treatment of soil enhances Verticillium infection of barberry and redbud, 996
---: storage, cold storage, urediospores of Puccinia graminis f. sp. tritici stored in liquid nitrogen at -196°C during 2 1/2 years showed no detectable change in virulence or color and maintained genetic stability, 695
Tenacity: abilities of 19 fungicides to be deposited and to persist at protective levels on plant leaves, Difolatan had the best ranking in the tests, 898
Tennessee: 625; 697
Texas: 72; 202; 1073
Thielavia basicola: 911
Thielaviopsis basicola: 238; 352
Tilia platyphyllos (see Lime-tree)
Tilletia caries: 544; 1132
--- contraversa: 613; 1132
--- foetida: 544
Timothy (Phleum pratense): occurrence in Wisconsin of halo blight of timothy incited by a Pseudomonas species closely related to P. coronafaciens, 361
Tobacco: black root rot, soil amendments and fungicide treatments controlled Thielaviopsis basicola in greenhouse tests, 352
---: Leveillula taurica found on tobacco in Iraq for the first time, 192
---: nematodes, effects of depth of fall tillage, selected hosts, and fallowing on the population density of Meloidogyne incognita and Pratylenchus zeae, 41
---: Verticillium wilt: benomyl applied to soil reduced loss of tobacco leaf caused by Verticillium wilt in New Zealand, 889
---: virus: tobacco ringspot virus, ash (Fraxinus americana) strain can be transmitted by dagger nematodes, Xiphinema americanum, 475; viruses of tobacco in North Carolina, 1013
Tobacco etch virus: on pepper, in North Carolina, 751; in Hawaii, 783
Tobacco mosaic virus: on pepper in North Carolina, 751; in Hawaii, 783
Tolyposporium penicillariae: 87
Tomato: anthracnose, caused by Colletotrichum coccodes, resistance in parents PI 272636 and 'Roma' appeared to be partially dominant with several genes involved, 253
---: bacterial wilt: grafting susceptible tomato scions to resistant rootstocks is effective method for control of bacterial wilt in Hawaii, 521; excised-petiole method of inoculation is a more sensitive method of detecting Corynebacterium michiganense from infected overwintered stems than cotyledonary inoculation, 654
---: crown gall, retardation after absorption of PB2-A bacteriophage by tomato plants 145
---: early blight, 30 varieties or breeding lines tested for comparative resistance, 807
---: Fusarium wilt: of tomato, Monograph No. 6, 653; Fusaria associated with wilted tomato plants in Massachusetts, 687
---: nematodes: mineral fertilizer and oat straw amendments reduced root knot, 1126; the number of Pratylenchus penetrans in soil was not correlated with the incidence of Verticillium wilt, 673
---: Verticillium wilt, controlled with several nematicides, benomyl and EDB, 26; no correlation found between number of P. penetrans in soil and incidence of wilt, 673
---: virus, inoculation of seedlings with tobacco mosaic virus by quick-dip method for screening large numbers of seedlings for susceptibility to strains of TMV, 375
Toxin: T-toxin from Race T of Helminthosporium maydis, as an indicator of resistance to southern corn leaf blight, 667

Transmission: rose spring dwarf is a graft-transmissible disease identified in 1968 in Rosa multiflora, 294; of pear virus by Psylla pyricola to cucumber, 295; of virus from Sphaerotheca lanestris and Erysiphe graminis to Chenopodium that resembles tobacco mosaic virus, 342; transmission of ash strain of tobacco ringspot virus by Xiphinema americanum, 475; of Helminthosporium maydis by infected corn seed, 486; of papaya leaf reduction virus by Myzus persicae, 526; of bean pod mottle virus to soybean by striped blister beetle, Epicauta vittata, 628; sap-transmissible viruses of elderberry in New York, 913; a sap-transmissible virus was isolated from mosaic-affected sorghum plants grown in southern France, 1090
Tree collapse: a serious problem of citrus in Izmir region of Turkey, cause unknown, 464; of citrus, from boron toxicity, 1132
Tree mortality: of loblolly pine in once-thinned plantations, from Fomes annosus, 625; of short leaf pine from F. annosus, estimated by aerial photography, 761; loss from F. annosus in shortleaf pine plantations in Missouri, 1085
Trichoderma sp.: 659
Tricothecium roseum: 283
Trillium grandiflorum: plants with a greening disorder of the flowers yielded mycoplasma-like bodies in phloem tissue, 1108
Tulip root: of oats, in California, tolerant variety, Curt is recommended where Ditylenchus dipsaci is a problem, 433
Turkey: 464; 1132
Turnip (See Crucifers)
Tylenchulus semipenetrans: 154
Typhula itoana: 1118

Uromyces phaseoli var. vignae: 384
Urophlyctis leproides: first confirmable occurrence of disease on sugarbeet in North America, in California, 30
Ustilago avenae: 1122
--- hordei: 544
--- nuda: 118; 1122
--- tritici: 1053
Utah: 1069

Valsa ambiens: 788
Variegation: of Trillium grandiflorum, the possibility proposed that T. grandiflorum var. variegatum should be regarded as a disease form and not with a varietal designation, 1108
Vectors associated with: barley yellow dwarf, Rhopalosiphum maidis and R. padi, 797; barley yellow dwarf, Schizaphis graminum, 874; bean pod mottle virus, transmitted by Epicauta vittata to soybean, 628; beet mosaic virus and beet western yellows virus, Myzus persicae, 676; birch bark disease, Cryptococcus fagi, 591; cereal enation disease of oats, by leafhopper, Cicadulina mbila, 802; corn stunt, Graminella nigrifrons, 273; curly-top virus in Iran, Circulifer opacipennis, and by C. tenellus in the U.S., 85; Dutch elm disease, Scolytus multistriatus, 271; Dutch elm disease, Scolytus multistriatus and Hylurgopinus rufipes, known distribution in Minnesota, 682; elm phloem necrosis, Scaphoideus luteolus, 1085; oak wilt by Pseudopityophthorus minutissimus, 108; 1106; pear decline, transmission of virus from pear to cucumber by Psylla pyricola, 295; X disease of peach, leafhopper, Scaphytopius acutus, 645
Venturia inaequalis: 828; 976
Verbena: ETMT fungicide superior to Dexon in control of pythiaceous fungi in bedding plants, 587
Verticillium albo-atrum: 26; 347; 569; 780; 898; 968; 977; 996
--- dahliae: 889; 916
--- malthousei: 120; 643; 684
Vetch: hairy, new record as host for lespedeza cyst nematode, 927
Viability: urediospores of Puccinia graminis sp. tritici were stored in liquid nitrogen for 10 years with little loss in viability, 871; of frozen conidia of apple scab fungus, Venturia inaequalis, 976
Vigna sinensis (see Cowpea)
Vinca: disease control of bedding plants with ETMT fungicide, 587; first noncucurbit host of wild cucumber mosaic virus, 546
Virginia: 122; 201; 419; 467
Virulence: comparative virulence of American and Australian isolates of Verticillium albo-atrum in cotton, 977
Virus: transmission of a pear virus to cucumber by Psylla pryicola, 295; Erysiphaceae transmit virus to Chenopodium, 342; virus-like symptoms found on a blackjack oak in Arkansas in form of mosaic, 411; of Colocasia and Xanthosoma in Puerto Rico, 506; hot water treatment of Prunus budsticks failed to eliminate 12 suspected viruses, 738; additional diagnostic host plants found for papaya mosaic virus and papaya ringspot, 785
---: alfalfa mosaic virus: a numerical system of differentiating strains of AMV, 127; alfalfa mosaic virus strains from carrot and parsley, in California, 322
---: ash strain, of tobacco ringspot virus, can be transmitted by dagger nematodes, 475
---: barley yellow dwarf: in Manitoba in 1970, 797; a new variant of barley yellow dwarf

transmitted specifically by Schizaphis graminum was found in New York, 874
---: bean pod mottle virus, transmission by Epicauta vittata to soybean, 628
---: beet mosaic virus (BMV) and beet western yellows virus (BWYV) of sugarbeets were most prevalent of aphid-borne viruses in Washington from 1967-1969, 103; occurrence of BMV and BWYV in Washington is related to occurrence of volunteer beets and also to the mode of vector transmission, 676
---: Bidens mottle virus, was isolated from naturally infected lettuce and endive in Florida, 1061
---: citrus: exocortis virus is associated with dwarfness in citrus trees on trifoliate orange, 998; greening disease of sweet orange, decreased with trunk injections of tetracyclines in South Africa, 747
---: corn stunt virus, johnsongrass found to be a host of a strain of corn stunt in South Carolina, 273
---: Eaton-bug, a disease of sweet cherry in Washington, 344
---: of elderberry, sap-transmissible viruses isolated in New York, 913
---: of geranium, plants differentiated from geranium anther callus were free from virus-like leaf symptoms present in the parent plants, 1017
---: papaya leaf reduction virus, transmission by Myzus persicae, 526
---: pea seedborne mosaic virus, over 600 cultivars and lines of Pisum sativum tested for resistance; two lines show promise, 408
---: peanut stunt virus, in Georgia, apparently poses no threat to peanut production at this time, 453
---: pear latent virus, Virginia crab apple as an indicator, 983
---: of pepper: tobacco etch virus (TEV) is most prevalent virus on pepper in North Carolina, 751; potato Y virus, tobacco mosaic, TEV and spotted wilt virus affect commercial and wild peppers in Hawaii, 783; potato virus Y, resistance to this virus in peppers is apparently influenced by environmental factors with summer resistance and susceptibility in winter, 604
---: rose: rose spring dwarf, a graft-transmissible disease of Rosa multiflora in California, care should be taken to avoid scionwood and rootstocks infected with the disease, 294; rose mosaic virus, use of Shirofugen flowering cherry as indicator for rose mosaic virus has limited reliability, 431; rose wilt virus, symptom differences between rose wilt virus and Verticillium wilt of roses, 916
---: of sorghum, mechanically transmissible virus found in France appears similar to johnsongrass strain of maize dwarf mosaic virus in the U.S. and sorghum red stripe virus in Italy, 1090
---: of strawberry, runner plants grown in Oregon in 1970 had lower levels of virus present than in 1969, 220
---: of sugarbeet: beet mosaic virus and beet western yellows virus in Washington, 103; 676; curly top virus in Iran, 85
---: sugarcane mosaic virus (SCMV), Krish sorghum (an interspecific hybrid) showed high resistance to four strains of SCMV, 596
---: of sweetpotato, found in 24 commercial clones and in 28 clones in the germ plasm collection in Puerto Rico, 902
---: tobacco: tobacco yellow dwarf, controlled in Australia with systemic biocides, with possible direct action on mycoplasma, 291; tobacco mosaic, a quick-dip technique for inoculation of tomato seedlings with TMV, 375; in North Carolina, TMV is most prevalent on flue-cured tobacco and tobacco etch virus most prevalent on burley tobacco, 1013
---: watermelon mosaic in squash, WMV-1 and WMV-2 found in New York and New Jersey, 132
---: western celery mosaic, occurrence of virus in umbelliferous weeds, 268; recovered from parsley and poison hemlock in California, 328
---: wild cucumber mosaic virus, found to occur in Marah oreganus (wild cucumber) in Oregon, 546
---: X disease of peach, eradication of infected chokecherry nearby appears the only practical means of control, 645

Washing: citrus, effect of time of washing and fungicide application on the decay of degreened oranges and tangerines, 149
Washington: 57; 103; 135; 234; 344; 514; 676; 730; 1056
Water: falling water drops as rain, or overhead irrigation is apparently the direct causal factor of flower blast of statice in Florida, 848
Water congestion: nonparasitic disease of pea, 249; of 321 pea lines tested, none was completely resistant to water congestion; the most tolerant strains were PI 166084 and PI 180329, 533
Water dip (see Temperature)
Wave years: of western gall rust infection on pine, 163
Weather: environmental conditions appeared to be major factors for occurrence of

downy mildew on peas, 730; climatic elements important as sugarbeet epidemiology indicators in Southwest for Cercospora leaf spot epidemics, 1073
Weed hosts: four weeds are hosts of Sclerotinia spp. in alfalfa fields, 1087
West Pakistan: 1130
West Virginia: 1106
Wheat: bunt: control of common bunt (Tilletia caries and T. foetida) and dwarf bunt (T. contraversa) with systemic fungicides, 1132; seed treatments with thiabendazole reduced incidence of dwarf bunt, 613; control of hill bunt (T. caries and T. foetida) of wheat in India with benomyl, 544
---: eyespot (Cercosporella), control of lodging due to eyespot with benomyl, 45
---: glume blotch, caused by Phoma insidiosa, not previously reported on wheat, 95
---: nematodes, Heterodera avenae invasion was not suppressed by benomyl, 67
---: rust: leaf: caused by Puccinia recondita, in North Dakota in 1970, 172; controlled with new systemic fungicide in experimental testing, 691
---: ---: stem: distribution, prevalence and new physiologic races of Puccinia graminis tritici in the U. S. in 1969, 187; a system of nomenclature for physiologic races, 718; summary of the races identified by area and source of collection in the USA in 1970, 986; an unrecognized source of overwintering wheat stem rust, 990
---: smut, loose smut, caused by Ustilago tritici, controlled with fungicides, 1053
White mold: of bean, caused by Sclerotinia sclerotiorum, controlled in field with fungicides, 234
White thread blight; of cacao, caused by two Marasmius species, 885
Wildrice (Zizania aquatica): stem rot, caused by Helminthosporium sigmoideum, isolated from cultivated wildrice plants in Minnesota, 498
Wilt: bacterial: of cucurbits, response of several cucurbit genera to Erwinia tracheiphila, 816; of tomato, control by grafting to resistant rootstocks, 521
---: Fusarium: estimated cotton loss by State in 1970, 504; of cotton, several varieties were found highly resistant over a 4-year period, 896; of pea, breakdown of resistance of pea plants to wilt by Pratylenchus penetrans, 1032; of petunia, controlled with methyl bromide, 580; of sunflower, 1124; of tomato, 20 isolates identified as to their pathogenicity to wilt-susceptible and wilt-resistant tomato, 687
---: oak wilt: Ceratocystis fagacearum is carried by oak bark beetles, 108; sampling methods, 281; three herbicides gave no control of mat production or bark beetle infestation of oaks, 1106
---: Verticillium: estimated loss of cotton from, 504; of cotton and olive, relation of strains of Verticillium albo-atrum to severity of wilt in cotton and olive in California, 780; of potato, suppressed with fumigation with 1,3-D + chloropicrin, 968; to tobacco, reduced with benomyl soil treatment, 889; of tomato, controlled with fumigants for up to 3 months, 26; of tomato, no correlation found between numbers of Pratylenchus in soil and incidence of tomato wilt, 673
Wind: as a possible important means of nematode distribution in West Texas, 61
Wisconsin: 249; 361; 533; 536; 648; 788
Witchweed (Striga lutea): five new fungi found to be pathogenic to witchweed, 861
Wood discoloration: of Katsura-tree, caused by Dothiorella ribis, 893
Wyoming: 336

Xanthomonas fragariae: 142
--- maculifoliigardeniae: 261
--- oryzae: 17; 22; 593
--- phaseoli: 505; 577; 609; 703
--- --- var. fuscans: 577; 703
--- pruni: 774
--- translucens f. sp. hordei: 365
--- --- f. sp. oryzicola: 404
Xanthosoma spp. (yautía): large percentage of yautía in Puerto Rico are infected with at least one virus, 506
Xiphinema americanum: 475

Yellow leaf disease: of arecanut (Areca cathecu), 170
Yellow poplar (Liriodendron tulipifera): Cylindrocladium floridanum caused severe mortality of seedling yellow-poplar in a tree nursery in North Carolina, 700
Yield loss: from southern corn leaf blight, from corn inbreds and hybrids with different cytoplasms, 115; of cotton, by parasitic diseases in 1970, 504

PLANT SCIENCE RESEARCH DIVISION

Prepared by Marguerite G. Benedict

☆U.S. GOVERNMENT PRINTING OFFICE: 1972-481-850/11